Lecture Notes in Microeconomics

(The Economic Agent) Ariel Rubinstein Solution Manual

for March 2011 version

This document is intended for use by instructors of courses in Microeconomics only. You have received it after agreeing not to transfer it to anyone else and that it is only for your personal use.

The document contains the solutions to all the questions that appear in the Jan 2010 version of the book. Be aware that the problem sets in the book will be modified in the future.

I would be grateful if you alert me to any mistakes you find in the manual.

My thanks to the many TAs who helped me create this manual and in particular to Benjamin Bachi who contributed so much to the current version.

Ariel Rubinstein rariel@post.tau.ac.il http://arielrubinstein.tau.ac.il

Problem Set 1 - Preferences

1. Let \succeq be a preference relation on a set *X*. Define $I(x)$ to be the set of all *y* ∈ *X* for which *y* \sim *x***. Show that the set (of sets!)** $\{I(x)|x \in X\}$ is a partition of *X***.**

(i) For all *x* and *y*, either $I(x) = I(y)$ or $I(x) \cap I(y) = \emptyset$.

Note that the indifference relation is symmetric and transitive.

Let $x, y \in X$, and assume that $I(x) \cap I(y) \neq \emptyset$, which means that there exists $z \in I(x) \cap I(y)$. Let *a* be any element in one of the sets, let us say $I(x)$. It means that *a* \sim *x*. But *z* \sim *x* and *z* \sim *y* since *z* \in *I*(*x*) \cap *I*(*y*). By symmetry of the indifference relation, $x \sim z$. By transitivity of \sim , it follows that $a \sim y$, and thus $a \in I(y)$.

(ii) For every $x \in X$, there is $y \in X$ such that $x \in I(y)$.

Let $x \in X$. The completeness of \geq means that $x \sim x$. Consequently, $x \in I(x)$.

2. Kreps (1990) introduces another formal definition for preferences. His primitive is a binary relation *P* **interpreted as "strictly preferred". He requires** *P* **to satisfy:**

Asymmetry: For no x, y do we have both xPy and yPx **.**

Negative Transitivity (NT): $\forall x, y, z \in X$, if xPy , then either xPz or zPy (or both). **Explain the sense in which Kreps' formalization is equivalent to the traditional definition.**

We will closely follow the proof presented in the lecture notes. The following steps are required:

a. Construct an interpretation-preserving function *T* that maps a binary relation *P* satisfying asymmetry and negative-transitivity into preference relations.

Consider the following candidate function *T*, which maps any relation *P* into a binary relation defined by

 $xT(P)y$ if not yPx .

Note that *T* preserves interpretation: If "*y* is not strictly preferred to *x*" according to Kreps' formalization, then "*x* is at least as good as *y*".

b. Prove that $T(P)$ is a preference relation.

Completeness of *TP*: Kreps' asymmetry property of *P* says that for any *x* and *y* in *X*, either not *xPy* or not *yPx*. Thus either $xT(P)y$ or $yT(P)x$.

Transitivity: Let $x, y, z \in X$ be such that $xT(P)y$ and $yT(P)z$. If not $xT(P)z$, then zPx . By *NT*, either yPx or zPy . Thus either not $xT(P)y$ or not $yT(P)z$, a contradiction.

c. Prove that *T* is one-to-one.

Let P_1 and P_2 be two different relations satisfying Kreps' properties. Then there is a pair *x*, *y* such that xP_1y and not xP_2y (or the opposite), and thus not $yT(P_1)x$ and *yT*(P_2)*x*. Thus, $T(P_1) \neq T(P_2)$, implying that *T* is one to one.

d. Prove that *T* maps onto all preference relations.

Let \succeq be a preference relation. Define *P*, by:

xPy if not $y \ge x$.

P preserves Kreps' properties:

Asymmetry: Since \succsim is complete, then we never have both xPy and yPx .

NT: Let $x, y, z \in X$ be such that xPy , and thus not $y \geq x$. Therefore it is not true that both $y \ge z$ and $z \ge x$. Therefore either zPy or xPz .

Finally, note that $T(P) = \succeq$.

3. Let *Z* **be a finite set and let** *X* **be the set of all nonempty subsets of** *Z***.** Let \succeq be a preference relation on X (not Z). Consider the following two **properties of preference relations on** *X***:**

1. If $A \geq B$ and C is a set disjoint to both A and B , then $A \cup C \geq B \cup C$, and **if** $A \succ B$ and C is a set disjoint to both A and B , then $A \cup C \succ B \cup C$.

2. If $x \in Z$ and $\{x\} \succ \{y\}$ $\forall y \in A$, then $A \cup \{x\} \succ A$, and **if** $x \in Z$ and $\{y\} \succ \{x\}$ $\forall y \in A$, then $A \succ A \cup \{x\}$.

a. Discuss the plausibility of the properties in the context of interpreting as the attitude of the individual toward sets from which he will have to make a choice at a "second stage".

Consider an appealing interpretation of the formal model: The elements in *Z* are the alternatives which might be chosen at the end of a decision process, a set *A* is a set of candidates to be considered seriously in the second stage. If we have in mind that the economic agent is certain about his preferences in the later stage then (a) is problematic: if the best element in menu *A* is better than the best in menu *B*, and menu *C* includes an even better element, then $A \geq B$ but $A \cup C \sim B \cup C$, violating (a). Also, if any element of menu *A* is strictly better than *z* ∈ *Z*, then A > $\{z\}$ but $A \sim A \cup \{z\}$, violating (b). The properties make more sense if the decision maker has in mind a tiny possibility that he will err in his choice or that there is a possibility that an alternative which he chooses will not be feasible at the end.

b. Provide an example of a preference relation that:

(1) Satisfies both properties.

The relation \geq defined by $A \geq B$ if $|A| \geq |B|$ satisfies (a) and (b) in a degenerate way (since for all x and y we have $\{x\} \sim \{y\}$).

A "better" class of examples (including the previous one): Let *X* be divided to two sets *G* and *B*. Define a preference relation by the utility function $u(A) = |A \cap G|$ − $|A \cap B|$. Clearly it satisfies both properties.

(2) Satisfies the first but not the second property.

Let *z*[∗] ∈ *Z*. Define \succeq over *X* whereby $A \succ B \Leftrightarrow z^* \in A, z^* \notin B$ and $A \sim B$ otherwise.

(a) Let $A, B, C \in X$ be such that $A \succeq B$ and *C* is disjoint to both *A* and *B*. If $A \succ B$, then $z^* \in A$ and $z^* \notin B$, C, which implies that $A \cup C \succ B \cup C$. If $A \sim B$, then either z^* is a member of both sets $A \cup C$ and $B \cup C$ or of none. In both cases $A \cup C \sim B \cup C$.

(not b) Let $A = \{z^*\}$ and $y \neq z^*$. Then $A \succ \{y\}$ but $A \sim A \cup \{y\}$ violating the second part of (b).

More generally, attach to each element x a non-negative number $v(x)$ and define a preference relation by a utility function $U(A) = \sum_{a \in A} v(a)$. Then the preference relation satisfies (a) but typically not (b) unless v gets only one positive and one negative values.

(3) Satisfies the second but not the first property.

Let \succeq^* be a preference relation on *Z*. Define \succeq by $A \succeq B$ if

(1) the \succcurlyeq^* –best element in A is strictly better than the \succcurlyeq^* –best element in B, or

(2) the agent is indifferent between the best elements but the \succ -worst element in A is weakly better than the \succeq^* –worst element in *B*.

(not a) Let $a \succ^* b \succ^* c \succ^* d$. Then $\{b\} \succ \{c\}$, but $\{b\} \cup \{a,d\} \sim \{c\} \cup \{a,d\}$. (b) Let $A \in X$ and $z \in Z$. If z is strictly \succeq^* -better (strictly \succeq^* -worse) then all $a \in A$, then $A \cup \{z\} > A \quad (A > A \cup \{z\})$.

c. Show that if there are $x, y, z \in Z$ such that $\{x\} \succ \{y\} \succ \{z\}$, then there is no **preference relation satisfying both properties.**

Assume \succsim satisfies (a) and (b), with $\{x\} \succ \{y\} \succ \{z\}$ for some $x, y, z \in Z$. From (b), $\{x\}$ > $\{x,y\}$ and $\{y,z\}$ > $\{z\}$. Applying (a) to the above, $\{x,z\}$ > $\{x,y,z\}$ and $\{x, y, z\}$ > $\{x, z\}$, a contradiction.

4. Let $>$ be an asymmetric binary relation on a finite set X that does not have cycles. Show (by induction on the size of X) that \ge can be extended to a **complete ordering.**

If a set *A* is finite and \geq is an acyclic relation on *A* (there are no cycles), then there must exist an $x \in A$ such that there is no $y \in A$ such that $y \succ x$.

Since *X* is finite, then there exists an $x_1 \in X$ such that there is no $y \in X$ such that *y* $> x_1$. Define $x_1 >^* y$ for all such $y \in X - \{x_1\}$. Again, there exists an $x_2 \in X - \{x_1\}$ such that there is no $y \in X - \{x_1\}$ such that $y \succ x_2$. Define $x_2 \succ^* y$ for all such y, and so on. By induction we can define \succ^* for all $x \in X$.

By construction, the relation \succ^* is complete, asymmetric, extends \succ and transitive: let $x_i \succ^* x_j$ and $x_i \succ^* x_h$. Then $i < j$ and $j < h$ and therefore, $x_i \succ^* x_h$.

5. Listen to the illusion called the Shepard Scale. (You can find it on the internet. Currently, it is available at http://asa.aip.org/demo27.html.) Can you think of any economic analogies?

The Shepard Scale consists of three separate scales that play the same tone at different octaves. As the notes ascend, one scale drops its pitch an octave, a change the listener does not notice because the other two scales continue to ascend monotonically, which "covers up" the drop. Several notes later, a second scale drops its pitch an octave, and so on. Thus, the Shepard Scale sounds as if it perpetually ascends, even though the same finite set of notes are repeated. See http://en.wikipedia.org/wiki/Shepard_tone for explanation.

The phenomenon is a reminder of an example due to Fishburn and LaValle (1988):

Many of us prefer the lottery f_2 to the lottery f_1 . However, one can easily construct the other 4 lotteries f_3 , f_4 , f_5 , f_6 , (increase the \$500 to \$1000 and reduce the other prizes by \$100), such that we would prefer f_3 to f_2, f_4 to f_3, \ldots and f_1 to f_6 .

See Peter C. Fishburn and Irving H. LaValle (1988). "Context-Dependent Choice with Nonlinear and Nontransitive Preferences", Econometrica, Vol. 56, 1221-1239. Stable URL: http://www.jstor.org/stable/1911365

Problem Set Two – Utility

1. The purpose of this problem is to make sure that you fully understand the basic concepts of utility representation and continuous preferences. Prove or disprove the following:

a. Is the statement "if both U and V represent \succeq , then there is a strictly **monotonic function** $f : \mathbb{R} \to \mathbb{R}$ such that $V(x) = f(U(x))$ " correct?

False: Let $X = \mathcal{R}$ and preferences be represented by the utility functions

$$
V(x) = x \quad \text{and} \quad U(x) = \begin{cases} x & \text{if } x \le 0 \\ x+1 & \text{if } x > 0. \end{cases}
$$

The only increasing function $f: \mathbb{R} \to \mathbb{R}$ that satisfies $V(x) = f(U(x))$ is

$$
f(x) = \begin{cases} x & \text{if } x \le 0 \\ 0 & \text{if } 0 < x \le 1 \\ x - 1 & \text{if } x > 1 \end{cases}
$$

which is not *strictly* increasing.

b. Can a continuous preference relation be represented by a discontinuous utility function?

True: The preferences $(x \ge y$ if $x \ge y$) is represented by U in (a) are continuous, though *U* is discontinuous.

c. Show that in the case of $X = \mathcal{R}$, the preference relation that is **represented by the discontinuous utility function** $u(x) = [x]$ **(the largest integer** *n* such that $x > n$ **)** is not a continuous relation.

 $1 > 1/2$, but $1 - \epsilon \sim 1/2$ for $\epsilon > 0$ small, violating C1.

d. Show that the two definitions of a continuous preference relation, C1 and C2, are equivalent to

1. Definition C3: ∀*x* ∈ *X***, the upper and lower contours** ${y \mid y \ge x}$ *and* ${z \mid x \ge z}$ are closed sets in *X***.**

Definition C4: $\forall x \in X$, the sets $\{y \mid y \succ x\}$ and $\{z \mid x \succ z\}$ are **open sets in** *X***.**

(C3 \Leftrightarrow C4) By completeness, the sets $\{y \mid x \succ y\}$ and $\{y \mid y \succ x\}$ are the complementary to $\{y \mid y \geq x\}$ and $\{y \mid x \geq y\}$ correspondingly. Thus the formers are open sets iff the later are closed sets.

(C1 \Rightarrow C4) Let $x \in X$ and $a \in \{y \mid y \succ x\}$. By C1, there exists an $\epsilon > 0$ such that $B_{\epsilon}(a) \subseteq \{y \mid y > x\}, (B_{\epsilon}(a)$ is the set of points in X that are less than ϵ distance from *a*). Thus $\{y \mid y \succ x\}$ is open. The argument for $\{z \mid x \succ z\}$ open is analogous.

(C4 \Rightarrow C1) Let us use the notation *B* \times *x* to mean that *y* \times *x* for all *y* \in *B*.

Assume first that there exists an $z \in X$ such that $x \succ z \succ y$. By C4, there exist $\epsilon_1, \epsilon_2 > 0$ such that $B_{\epsilon_1}(x) > z$ and $z > B_{\epsilon_2}(y)$. Let $\epsilon = \min\{\epsilon_1, \epsilon_2\}$. By transitivity,

every point in $B_{\epsilon}(x)$ is strictly better than every point in $B_{\epsilon}(y)$.

Next, assume that there does not exist a $z \in X$ such that $x \succ z \succ y$. As above, by C4 there exists an $\epsilon > 0$ such that $B_{\epsilon}(x) > y$ and $x > B_{\epsilon}(y)$. Since there is no *z* such that $x \succ z \succ y$, then $B_{\epsilon}(x) \succ z$ and $y \succ z$ $B_{\epsilon}(y)$, and thus by transitivity, every point in $B_{\epsilon}(x)$ is strictly better than every point in $B_{\epsilon}(y)$.

2. Give an example of preferences over a countable set in which the preferences cannot be represented by a utility function that returns only integers as values.

Let $X = N$, which is countable. Define preferences to be such that

 $1 \succ 3 \succ 5 \succ \ldots \succ 2 \succ 4 \succ \ldots$

By contradiction, assume that there exists a utility function $u: X \rightarrow \mathbb{Z}$ that represents \geq . Then $u(1) = N$ and $u(2) = n$ for some $n, N \in \mathbb{Z}$. But there are an infinite number of odd numbers, implying that *u* maps to an infinite number of integers between *n* and *N*, a contradiction.

3. Let \succcurlyeq be continuous preferences on a set $X \subseteq \mathbb{R}^n$ which contains the **interval connecting the points** *x* **and** *z*. **Show that if** $y \in X$ **and** $x \ge y \ge z$, **then there is a point** *m* on the interval connecting *x* and *z* such that $y \sim m$.

Construct inductively the sequence $\{(x^n, z^n, m^n)\}$ as follows: Start with define $x^0 = x$, $z^0 = z$ and the midpoint $m^0 = 1/2x^0 + 1/2z^0$.

If $m^n \sim y$ then we found the point we look for. Otherwise, $m^n > y$ or $y > m^n$.

If $m^n > y$ let $x^{n+1} = m^n$ and $z^{n+1} = z^n$.

If $y > m^n$ let $x^{n+1} = x^n$ and $z^{n+1} = m^n$.

In any case define $m^{n+1} = 1/2x^{n+1} + 1/2z^{n+1}$.

If none of the points $m^n \sim y$ then $x^n \succ y \succ z^n$ for all *n*. Both sequences (x^n) and $(zⁿ)$ converge to some $m[*]$ on the interval between x and z.

Since \ge is continuous, then $m^* \ge y$ and $y \ge m^*$, and thus $m^* \sim y$.

4. Consider the sequence of preference relations $(\geq^n)_{n=1,2,\ldots}$, defined on \mathfrak{R}^2_+ where \succcurlyeq^n is represented by the utility function $u_n(x_1, x_2) = x_1^n + x_2^n$. We will say that the sequence \succsim ^{*n*} converges to the preferences \succsim ^{*} if for every *x* and *y* such that $x >^* y$, there is an *N* such that for every $n > N$ we have $x >^n y$. **Show that the sequence of preference relations** \geq^n **converges to the preferences** \succeq^* **which are represented by the function** $max\{x_1, x_2\}$.

Let $x \succ^* y$. Since $\max\{x_1, x_2\} > \max\{y_1, y_2\}$, then there exists an $\epsilon > 0$ such that $\max\{x_1, x_2\}$ > $(1 + \epsilon) \max\{y_1, y_2\}$. Consequently, for *n* large enough $[\max\{x_1, x_2\}]^n > 2[\max\{y_1, y_2\}]^n$. But $x_1^n + x_2^n \geq [\max\{x_1, x_2\}]^n$ and $2 \max \{y_1, y_2\}^n \geq y_1^n + y_2^n$, and thus $x >^n y$ for *n* large enough.

5. The following is a typical example of a utility representation theorem: Let X = \mathfrak{R}_+^2 . Assume that a preference relation \succsim satisfies the following three **properties:**

ADD: $(a_1, a_2) \ge (b_1, b_2)$ implies that $(a_1 + t, a_2 + s) \ge (b_1 + t, b_2 + s) \; \forall s, t$. **SMON:** If $a_1 \geq b_1$ and $a_2 \geq b_2$, then $(a_1, a_2) \geq (b_1, b_2)$. In addition, if either

 $a_1 > b_1$ or $a_2 > b_2$ then $(a_1, a_2) > (b_1, b_2)$.

CON: Continuity.

a. Show that if has a linear representation (that is, are represented by a utility function $u(x_1, x_2) = \alpha x_1 + \beta x_2$ with $\alpha, \beta > 0$), then \succeq satisfies ADD, **SMON, CON.**

ADD: Let $s, t \in \mathcal{R}$ and $x, y \in X$ be such that $x \geq y$. Note that $(x_1, x_2) \ge (y_1, y_2) \Leftrightarrow \alpha x_1 + \beta x_2 \ge \alpha y_1 + \beta y_2 \Leftrightarrow \alpha (x_1 + t) + \beta (x_2 + s) \ge \alpha (y_1 + t) + \beta (y_2 + s)$ $u(x_1 + t, x_2 + s) \ge u(y_1 + t, y_2 + s) \Leftrightarrow (x_1 + t, x_2 + s) \ge (y_1 + t, y_2 + s).$

SMON: Let $x, y \in X$ be such that $x_1 \ge y_1$ and $x_2 \ge y_2$ with at least one strict inequality. Since α , $\beta > 0$, then $\alpha x_1 + \beta x_2 > \alpha y_1 + \beta y_2$, which implies that

 (x_1, x_2) \succ (y_1, y_2) .

CON: $u(x_1, x_2)$ is continuous, and thus \succeq is continuous.

b. **Show that for any pair of the three properties there is a preference relation that does not satisfy the third property.**

Satisfies only ADD, SMON: Lexicographic preferences satisfy *ADD* and *SMON*, but are not continuous (see the lecture notes).

Satisfies only ADD, CON: The preferences represented by $u(x_1, x_2) = x_1 - x_2$ satisfy *ADD* and *CON*, but not *SMON* since $(1,1) \succ (1,2)$.

Satisfies only MON, CON: Preferences represented by $u(x_1, x_2) = x_1^2 + x_2^2$ satisfy *SMON* and *CON*, but not *ADD* since $(3,0) > (2,1)$ and $(3,3) < (2,4)$.

c. Show that if satisfies the three properties, then it has a linear representation.

Assume first that x and y are two different points such that $x \sim y$. Then:

(i) $(x + y)/2 \sim y$. Otherwise, $(x + y)/2 \succ y$ would imply that

 $x = \frac{x+y}{2} + \frac{x-y}{2} > y + \frac{x-y}{2} = \frac{x+y}{2} > y$. by ADD, a contradiction.

(ii) $z = (1 - \alpha)x + \alpha y \sim x$ for $\alpha \in [0, 1]$. Define $\{(x^n, y^n)\}$ inductively as follows: let $x^0 = x$, $y^0 = y$. Let $m^0 = (x^0 + y^0)/2$.

Assume *z* belongs to $[x^n, y^n]$ and its length is $1/2^n$ the length of $[x, y]$. The point *z* belongs to at least one of the intervals $[x^n, m^n]$ and $[m^n, y^n]$. Define $[x^{n+1}, y^{n+1}]$ to be one of those intervals which contains z . Now, all $x^n \sim x$ for all n . The sequence $x^n \rightarrow z$, therefore by continuity $z \sim x$.

(iii) Let *z* be on the line which connects *x* and *y*, $z \sim x$. Without loss of generality, assume that *z* is closer to *x*. There is *n* such that $w = z + n(y - x)$ is between *x* and *y*. By ADD if $a - x = b - y$ (that is a-b=x-y) then $a \sim b$. Thus by transitivity $z \sim w \sim x$.

By SMON there is an $\varepsilon > 0$ such that $a = (x_1 + \varepsilon, x_2) > x > (x_1, x_2 - \varepsilon) = b$. By question 3, there exists *y* (different than *x*) on the interval which connects *a* and *b* such that $x \sim z$. Thus, every point is on a difference line which is a line. The indifference lines must be parallel since otherwise we will get a contradiction to *ADD*.

d. Characterize the preference relations which satisfy ADD, SMON and an additional property MUL:

 $(a_1, a_2) \geq (b_1, b_2)$ *implies that* $(\lambda a_1, \lambda a_2) \geq (\lambda b_1, \lambda b_2)$ *for any* $\lambda \geq 0$ *.*

Define $s = \sup\{x | (0,1) \succ (x,0)\}$ (by SMON the set is not empty).

Case (1): $s = \infty$ or $s = 0$: the preferences must be lexicographic with priority for the second or first components, respectively.

Assume $s = \infty$.

If $a_2 > b_2$ then $(a_1, a_2) > (b_1, b_2)$ iff $(a_1, a_2 - b_2) > (b_1, 0)$ (by ADD) iff

 $(a_1/(a_2 - b_2), 1)$ > $(b_1/(a_2 - b_2), 0)$ (by MUL), which is always true (by $s = \infty$). If $a_2 = b_2$ then $(a_1, a_2) \succ (b_1, b_2)$ iff $a_1 > b_1$ (by SMON).

Thus, we have a lexicographic relation with priority for the second component.

If $s = 0$ then it follows that $s = \sup\{y | (1,0) \succ (0,y)\} = \infty$ and the preferences must be lexicographic with priority for the first component.

Case (2): $\infty > s > 0$

Let (a_1, a_2) and (b_1, b_2) be two vectors with $a_1 \leq b_1$. (a_1, a_2) relates to (b_1, b_2) as $(0, a_2 - b_2)$ relates to $(b_1 - a_1, 0)$ (by ADD) and thus as $((b_1 - a_1)/(a_2 - b_2), 0)$ relates to $(0,1)$. This relation is determined by the comparison of $(b_1 - a_1)/(a_2 - b_2)$ to *s*, which is equivalent to the comparison of $a_1 + sa_2$ and $b_1 + sb_2$.

Therefore, if $(0,1) \sim (s,0)$ then $x_1 + sx_2$ represents the preferences. If $(0,1)$ > $(s,0)$ or $(0,1)$ < $(s,0)$ then the preferences are lexicographic with the first priority to $x_1 + sx_2$ and the second to x_2 or x_1 accordingly.

6. Let *X* be a finite set and let (\succcurlyeq, \succ) be a pair where \succcurlyeq is a preference **relation and is a transitive sub-relation of (by sub-relation, we mean** $x \rightarrow y$ implies $x \rightarrow y$). We can think about the pair as representing the **responses to the questionnaire** *A* where $A(x, y)$ is the question:

How do you compare *x* **and** *y***? Tick one of the following five options:**

 \Box **I** very much prefer *x* over $y(x \rightarrow y)$

 \Box **l** prefer *x* over y ($x > y$)

I am indifferent (*I***)**

 \Box **l** prefer *y* over x ($y > x$)

 \Box **I** very much prefer *y* over x ($y \gg x$)

Assume that the pair satisfies extended transitivity: If $x \gg y$ and $y \gtrsim z$, or if $x \geq y$ and $y \gg z$ then $x \gg z$. We say that a pair (\geq, \gg) is represented by a **function** *u* **if**

 $u(x) = u(y)$ iff $x \sim y$,

 $u(x) - u(y) > 0$ iff $x > y$, and

$$
u(x) - u(y) > 1
$$
 iff $x \succ y$.

Show that every extended preference $(\succcurlyeq, \succcurlyeq)$ is represented by a function *u***.**

Denote $A \succ B$ if $a \succ b$ for all $a \in A$ and $b \in B$. Let $X_1, X_2, ..., X_K$ be the \succeq indifference sets such that $X_K > X_{K-1} > ... > X_1$. Define first $u(X_1) = 0$.

Let us define $u(X_k)$ for $k > 1$.

(1) if $X_k \gg X_{k-1}$, then $u(X_k) = u(X_{k-1}) + 2$

(2) if X_k is not >> even of X_1 , then $u(X_k) \in (u(X_{k-1}),1)$

(3) otherwise, there exists a maximal $m(k)$ such that $X_k \gg X_{m(k)}$. Define $u(X_k)$

such that $u(X_k) > u(X_{k-1})$ and $1 + u(X_{m(k)+1}) > u(X_k) > u(X_{m(k)}) + 1$.

Clearly, $x \sim y$ iff $u(x) = u(y)$

Also, if $x > y$ then $u(x) > u(y)$, since we picked $u(X_k)$ as an increasing sequence. Finally, if $x \rightarrow y$, $x \in X_k$ and $y \in X_m$ then $m(k) \ge m$ and

 $u(x) > u(X_{m(k)}) + 1 \ge u(y) + 1.$

7. Utility is a numerical representation of preferences. One can think about the numerical representation of other abstract concepts. Here, you will try to come up with a possible numerical representation of the concept "approximately the same" (see Luce (1956) and Rubinstein (1988)). For simplicity, let $X = [0,1]$. Consider the following six properties of S :....

We will focus here only on (c).

Let *S* **be a binary relation that satisfies the above six properties and let** $\epsilon > 0$. Show that there is a strictly increasing and continuous function *H* : *X* → \Re such that aSb \Leftrightarrow $|H(a) - H(b)| \le \epsilon$.

Note the definitions of $m(x)$ and $M(x)$ in the question.

Define $\{x_n\}$ by $x_0 = 0$, $x_1 = M(0)$, $x_2 = M(x_1) = M(M(0))$ and so on. By S6, $\{x_n\}$ is increasing and bounded above by 1, and thus $\{x_n\}$ converges to $x^* \leq 1$. By S5, there exists an *N* such that $x_{N-1}Sx^*$, and thus $x^* \leq M(x_{N-1}) = x_N$. Since x^* is the upper bound of $\{x_n\}$, then $x^* = 1$ by S6. Define *N* to be the smallest integer such that $x_N = 1$, and thus $0 = x_0 < ... < x_N = 1$.

Lemma 1: If $a \in [x_n, x_{n+1}]$, where $1 \le n \le N-1$, then $m(a) \in [x_{n-1}, x_n]$. Proof: Since $x_n S x_{n+1}$, then $x_n S a$ by S4, and thus $m(a) \leq x_n$. Moreover,

 $x_{n-1} \leq m(a)$, as otherwise $m(a) < x_{n-1}$ and $M(x_{n-1}) = x_n \leq M(m(a))$, violating the assumption that *M* increasing.

Lemma 2: $m(a)$ is strictly increasing and continuous on $(x_1, 1]$.

Proof: $m(a) > 0$ if $a > x_1$, as otherwise aS_0 , and thus $M(0) \ge a > x_1$, a contradiction. By S6, the lemma is proved.

Define

$$
H(a) = \begin{cases} \frac{\epsilon}{x_1} a & \text{if } a \leq x_1 \\ H(m(a)) + \epsilon & \text{if } a > x_1. \end{cases}
$$

H is clearly continuous and strictly increasing on $[0, x_1]$, with $H(x_1) = \epsilon$.

If $a \in (x_1, x_2]$, then $H(a) = \epsilon[m(a)/x_1 + 1]$ since $m(a) \in [0, x_1]$ by Lemma 1. Thus *H* is strictly increasing and continuous on $(x_1, x_2]$ by Lemma 2. Since $m(x_1) = 0$, then $H(x) \to \epsilon$ as $x \to x_1$ from the right, and thus *H* in continuous and strictly increasing on $[0, x_2]$, with $H(x_2) = 2\epsilon$.

More generally, if $a \in (x_n, x_{n+1}]$, where $n \leq N-1$, then $m(a) \in [x_{n-1}, x_n]$, $m(m(a)) \in [x_{n-2}, x_{n-1}]$ and so on by Lemma 1. Therefore $H(a) = \epsilon[m(\dots m(a), \dots)/x_1 + n]$, which is strictly increasing and continuous by Lemma 2, where $m(...m(a)...)$ applies *m* inductively *n* times. Since $H(x) \rightarrow n\epsilon$ as $x \rightarrow x_n$ from the right, then *H* is strictly increasing and continuous on $[0, x_{n+1}]$.

Let $a, b \in [0, 1]$ where $a < b$. If $b \le x_1$, then *H* represents *S* by S4. Otherwise, aSb iff $H(m(b))$ ≤ $H(a)$ < $H(b)$ iff $|H(b) - H(a)|$ ≤ ϵ , where the first iff follows from *aSb* iff $m(b) \le a < b$ and *H* strictly increasing, and the second iff follows from $H(b) = H(m(b)) + \epsilon.$

Problem Set 3 – Choice

Problem 1:

The following are descriptions of decision making procedures. Discuss whether the procedures can be described in the framework of the choice model discussed in this lecture and whether they are compatible with the "rational man" paradigm.

a. The decision maker (DM) has in mind a ranking of all alternatives and chooses the alternative that is the worst according to this ranking.

Let \geq be the preferences reflecting the DM's ranking. The choice procedure is rationalized by the preference relation \succcurlyeq^* , where $b \succcurlyeq^* a$ whenever $a \succcurlyeq b$.

b. The DM chooses an alternative in order to maximize another person's suffering.

Assuming that the relation "the other person suffers more from *x* than he does from *y*" is complete and transitive, the DM is maximizing a well-defined preference relation.

c. The DM asks his two children to rank the alternatives and then chooses the alternative that is the best "on average".

The question is, of course, what does the expression "on average" mean. If the DM ranks all alternatives in *X* and uses the ranking to attach the number to each alternative *a* in any set *A* (independently of *A*, then the DM's behavior is consistent with rationality. But if the score of an alternative is recalculated for every choice set then his behavior may be inconsistent with the rational man paradigm. For example, assume that one child ranks the alternatives *a*,*d*, *e*,*b*, *c* and the other as *b*, *c*, *a*, *d*, *e*. Then, the element *a* is chosen from the set $\{a, b, c, d, e\}$ while *b* is chosen from $\{a, b, c\}$.

d. The DM has an ideal point in mind and chooses the alternative that is closest to it.

Let x be the ideal point and $d(a, b)$ the distance function between $a, b \in X$. The behavior is rationalized by the preferences represented by $u(a) = -d(a,x)$.

e. The DM looks for the alternative that appears most often in the choice set.

A choice function *C* is not well-defined. The DM's behavior is different when faced with the group of elements (a, a, b) than when faced with the group (a, b, b) . even though in both cases he chooses from the set $\{a, b\}$.

f. The DM always selects the first alternative that comes to his attention.

C is not well-defined: $C(< a,b>) = \{a\} \neq \{b\} = C(< b,a>)$, although the choice set in the two cases is the same.

g. The DM has an ordering in mind and always chooses the median element.

C violates condition α . Assume that the order of the grand set $X = \{a, b, c, d, e\}$ is alphabetical. Then, $C({a,b,c,d,e}) = {c}$ but $C({a,b,c}) = {b}.$

Problem 2:

Let's say that you are to make a choice from a set *A***. Consider two procedures:**

(a) You choose from the entire set or (b) You first partition *A* **into the subsets** *A*¹ **and** *A*2**, then make a selection from each of the subsets and finally make a choice from the two selected elements**.

a. Formulate a "path independence" property (for single-valued choice functions).

Let A_1 and A_2 be a partition of A. By partition, we mean $A_1, A_2 \neq \emptyset$, $A = A_1 \cup A_2$ and $A_1 \cap A_2 = \emptyset$. *C* satisfies path independence if

$$
C(A) = C({C(A1), C(A2)}).
$$

b. Show that the rational decision maker satisfies this property.

Let A_1 , A_2 be a partition of A . Let $x = C_1(A)$ and $x \in A_i$. Then $x = C_1(A_i)$ and x is also the best choice from $\{C_*(A_1), C_*(A_2)\}.$

c. Find examples of choice procedures that do not satisfy this property.

(i) The "second best" procedure. If $x > y > z$, then $C({x, y, z}) = y$ while $C({C(\{x,y\},C(\{z\})}) = \{z\}.$

(ii) Let *X* be partitioned into *Y* and *Z* and let \ge be an ordering on *X*. Let *C*(*A*) be the \triangleright -minimal element if all alternatives in *A* are in *Y* and the \triangleright -maximal alternative otherwise. If $Y = \{a,b\}$, $Z = \{c,d\}$ and $a \succ b \succ c \succ d$, then $\{C(\{a,b,c,d\}) = a$ but $C({C(\{a,b\}), C({\{c,d\}})) = b$.

d. Show that if a choice function satisfies path independence, then it is consistent with rationality.

We will show that Condition α is satisfied. Let $A \subset B \subseteq X$ be such that $C(B) \in A$. By path independence, $C(B) = C({C(A), C(B \setminus A)}$. Since $C(B)$ is in *A*, then it is not in *B* \setminus *A*. Therefore *C*(*B*) is identical to *C*(*A*).

e. Find an example of a multi-valued choice function satisfying path independence which cannot be rationalized.

In the context of choice correspondences, path independence implies:

$$
C(A) = C(C(A_1) \cup C(A_2)).
$$

Let $a > b > c > d$ and $C(A)$ be a set which contains the best and worst elements in *A*. *C* satisfies path independence (verify) but violates WA since $a \in C(\{a, b, c, d\})$ and $c \in C({a,b,c})$ but $c \notin C({a,b,c,d})$.

Problem 3:

Let *X* **be a finite set. Check whether the following three choice correspondences satisfy WA:**

1) $C(A) = \{x \in A \mid \text{the number of } y \in X \text{ for which } V(x) \geq V(y) \text{ is at least } }$ $|X|/2$ and if this set is empty, then $C(A) = A$.

By *C*, a satisfactory element is one which is in the upper half of the elements in the grand set.

C satisfies *WA*. Define:

G = $\{x \in X \mid V(x) \geq V(y) \}$ for at least $|X|/2$ alternatives $y \in X\}$.

Let $x, y \in A \cap B$ such that $x \in C(A)$ and $y \in C(B)$.

If $x \in G$, then $x \in C(B)$.

If $x \notin G$, then there are no elements of *A* in *G* and thus $y \notin G$. Since $y \in C(B)$, then $C(B) = B$ and thus $x \in C(B)$ as well.

Alternatively: define $u(x) = 1$ if $x \in G$ and 0 otherwise. Clearly, $C_u = C$.

2) $D(A) = \{x \in A \mid \text{the number of } y \in A \text{ for which } V(x) \geq V(y) \text{ is at least }$ |*A*|/2**.**

By *D*, a satisfactory element is one in the upper half of the elements of the choice set.

It is not necessarily consistent with the rational man paradigm: Let $V(a) > V(b) > V(c) > V(d) > V(e)$. Then, $c \in C({a,b,c,d,e})$ and $a \in C({a,b,c})$, but $c \notin C({a,b,c}).$

3) $E(A) = \{x \in A \mid x \succeq_1 y \text{ for every } y \in A \text{ or } x \succeq_2 y \text{ for every } y \in A\}$, where ≥ 1 and ≥ 2 are two orderings over X.

By *E*, a satisfactory element is one which is optimal according to one of the two criteria. It is not necessarily consistent with the rational man paradigm. Let $x \succ_1 y \succ_1 z$ and $y \succ_2 z \succ_2 x$. Then, $z \in C(\{x, z\})$ and $x \in C(\{x, y, z\})$, but $z \notin C(\{x,y,z\})$.

Problem 4:

Consider the following choice procedure: A decision maker has a strict ordering \succeq over the set *X* and assigns to each $x \in X$ a natural number $class(x)$ **to be interpreted as the "class" of** *x***. Given a choice problem** *A*, **he chooses the best element in** *A* **from those belonging to the most common class in A (i.e., the class that appears in** *A* **most often). If there is more than one most common class, he picks the best element from the members of** *A* **that belong to a most common class with the highest class number.**

a) Is this procedure consistent with the "rational man" paradigm?

No. Let $a > b > c > d > e$, $class(a) = class(b) = class(c) = 1$ and $class(d) = class(e) = 2.$

 C { $\{a, b, c, d, e\}$ } = *a* but C { $\{a, d, e\}$ } = *d*, thus violating α .

b) Define the relation xPy if x is chosen from $\{x, y\}$. Show that the relation *P* **is a strict ordering (complete, asymmetric and transitive).**

By definition, P is complete and asymmetric. We will see that it is also transitive. That is, if *x Py* and *yPz*, then *xPz*.

If xPy and yPz , then $\lceil class(x) \rangle$ *class* (y) or $class(x) = class(y)$ and $x \ge y$, and $[class(y) > class(z)$ or $class(y) = class(z)$ and $y \ge z$. If either $class(x) > class(y)$ or $class(y) > class(z)$, then $class(x) > class(z)$ and $C(\{x,z\}) = \{x\}$. Otherwise, $class(x) = class(z)$ and $x \ge z$ and thus $x \in C(\{x, z\})$.

Alternatively: Note that *P* is identical to the lexicographic preferences with first priority given to class and second priority to the relation \succeq .

Problem 5:

Consider the following two choice procedures. Explain each procedure and try to persuade a skeptic that they "make sense". Determine For each of them whether they are consistent with the "rational man" model.

a) The primitives of the procedure are two numerical (one-to-one) functions *u* **and** *v* **defined on** *X* **and a number** *v*[∗]**. For any given choice problem** *A***, let** *a*[∗] ∈ *A* **be the maximizer of** *u* **over** *A* **and let** *b*[∗] ∈ *A* **be the maximizer of** ν **over** A. The decision maker chooses a^* if $\nu(a^*) \geq \nu^*$ and b^* if $\nu(a^*) < \nu^*$.

One interpretation of this procedure is that the DM actually wants to maximize *v* but pretends to maximize *u*. If the maximization of *u* yields a result which is too bad for him, he abandons the pretense and maximizes *v*. The procedure may fail condition α . For example,

Then $C({x, y, z}) = z$ but $C({y, z}) = y$.

b) The primitives of the procedure are two numerical (one-to-one) functions *u* **and** *v* **defined on** *X* **and a number** *u*[∗]**. For any given choice problem** *A*, the decision maker chooses the element $a^* \in A$ that maximizes u **if** $u(a^*) \geq u^*$ and the element $b^* \in A$ that maximizes v if $u(a^*) < u^*$.

In this case, the DM, cares about the value of *u* only if it is at least *u*[∗]. Otherwise, he cares about *v*. The DM behaves as if he is maximizing lexicographic preferences with first priority given to the function $u'(x)$, which receives the value *u*(*x*) if *u*(*x*) ≥ *u*[∗] and *u*[∗] − 1 otherwise, and second priority to *v*(*x*).

Problem 6:

The standard economic model assumes that choice is made from a set. Let us construct a model where the choice is assumed to be made from a list (note that the list $\langle a, b \rangle$ is distinct from $\langle a, a, b \rangle$ and $\langle b, a \rangle$). Let *X* be a finite **grand set. A list is a nonempty, finite vector of elements in** *X***. In this problem, consider a choice function** *C* **to be a function that assigns a single element from** $\{a_1, \ldots, a_k\}$ to each vector $L = \langle a_1, \ldots, a_k \rangle$. Let $\langle L_1, \ldots, L_m \rangle$ be the concatenation of the *m* lists L_1, \ldots, L_m (note that if the length of L_i is k_i , then the length of the concatenation is $\sum_{i=1,...,m} k_i$). We say that L^* extends the list

L if there is a list *M* such that $L' = \langle L, M \rangle$.

We say that a choice function C satisfies property I if for all L_1, \ldots, L_m ,

 $C(\langle L_1, \ldots, L_m \rangle) = C(\langle C(L_1), \ldots, C(L_m) \rangle).$

a) Interpret Property *I***. Give two examples of choice functions that satisfy** *I* **and two examples that do not.**

Property *I* is analogous to path independence.

Two choice functions that satisfy *I*:

(i) Choose the first alternative in *L*.

(ii) Choose the first alternative in *L* that is "at least as good as" some $\tilde{x} \in X$ and choose the last element in *L* if there is no such alternative.

Two choice functions that violate *I*:

(i) Choose the second alternative in *L*.

(ii) Choose the last alternative such that the alteratives from the start of the sequence up to that alternative are in ascending order.

b) Define formally the following two properties of a choice function:

Order Invariance OI: A change in the order of the elements in the list does not alter the choice.

Let $L = \langle a_1, \ldots, a_K \rangle$. A permutation of *L* is a list $L^{\pi} = \langle a_{\pi(1)}, \ldots, a_{\pi(K)} \rangle$, where π is a permutation of $\{1, \ldots, K\}$. *C* satisfies OI if $C(L) = C(L^{\pi})$ for every permutation π .

Duplication Invariance DI: Deleting an element that appears elsewhere in the list does not change the choice.

C satisfies DI if $C(L) = C(L')$ whenever $L = \langle \langle L_1 \rangle, x, \langle L_2 \rangle \rangle$, $L' = \langle \langle L_1 \rangle, \langle L_2 \rangle \rangle$ and *x* appears in either L_1 or L_2 .

c) Characterize the choice functions that satisfy the following three properties together: Order Invariance, Duplication Invariance and Property *I***.**

Claim: Let *C* be a choice function over the *lists* of *X*. If *C* satisfies OI, DI and I, then there exists a rationalizable choice function *C* over the *sets* of *X* such that $C(L) = \overline{C}(\langle L \rangle)$, where $\langle L \rangle$ is the *set of elements* in *L*.

Proof: Let *K*,*L* be two lists such that $\{K\} = \{L\}$. By OI and DI, choice is preserved when the duplicate alternatives in both lists are removed and the resulting lists are reshuffled so that the remaining alternatives appear in the same order. Thus, $C(K) = C(L)$ and thus \overline{C} is well-defined and single-valued.

By Problem 2(d), showing that \overline{C} satisfies path independence is sufficient for

rationalizability. For any set $S \subseteq X$, define $\langle S \rangle$ to be some *list* of elements in *S*. Let $A, B \subseteq X$ be disjoint. Then:

$$
\overline{C}(A \cup B) = C(\langle \langle A \rangle, \langle B \rangle \rangle) \qquad \text{by def. of } \overline{C}
$$
\n
$$
= C(\langle C(\langle A \rangle), C(\langle B \rangle) \rangle) \text{ by property } I
$$
\n
$$
= C(\langle \overline{C}(A), \overline{C}(B) \rangle) \text{ by def. of } \overline{C}
$$
\n
$$
= \overline{C}(\langle \overline{C}(A), \overline{C}(B) \rangle) \text{ by def. of } \overline{C}
$$

Assume now that at the back of the decision maker's mind there is a value function *u* defined on the set *X* (such that $u(x) \neq u(y)$ for all $x \neq y$). For any **choice function** *C*, define $v_c(L) = u(C(L))$. We say that *C* accommodates a longer list if whenever L' extends L , $v_C(L') \ge v_C(L)$ and there is a pair of lists L' and L , such that L' extends L and $v_{C}(L') > v_{C}(L)$.

d) Give two interesting examples of choice functions that accommodate a longer list.

(i) Choose the *u*-maximal element in *L*.

(ii) Choose the second *u*-best alternative in *L*.

e) Give two interesting examples of choice functions which satisfy property *I* **but do not accommodate a longer list.**

(i) Choose the first alternative in *L* that yields at least utility *ũ* and choose the last alternative in *L* if $u(x) < \tilde{u}$ for all $x \in L$.

(ii) Choose the first element in *L*.

Problem 7:

We say that a choice function *c* **is lexicographically rational if there exists a profile of preference relations** $\{≻_a \}$ _{*a∈X*} (not necessarily distinct) and an **ordering** O **over** X **such that for every set** $A \subset X$, $c(A)$ is the \succ_a -maximal **element in** *A***, where** *a* **is the** *O* **-maximal element in** *A***.**

A decision maker who follows this procedure is attracted by the most notable element in the set (as described by *O***). If** *a* **is that element, he applies** the ordering a_a and chooses the a_a -best element in the set.

We say that *c* **satisfies the reference point property if for every set** *A***, there exists** $a \in A$ such that if $a \in A'' \subset A' \subset A$ and $c(A') \in A''$, then $c(A'') = c(A')$.

a. Show that a choice function *c* **is lexicographically rational if and only if it satisfies the reference point property.**

 Assume *c* is lex. rational. For every set *A* we'll show that the *O* − *maximal* element $a \in A$ satisfies the requirement of the reference point property. Note that for any A' ⊂ A containing a , a is still the O − $maximal$ element. Thus, for all subsets of of *A* containing a , $c(A')$ is determined by \succ_a . Thus condition α will be satisfied for all subsets of *A* containing *a*. This gives us the reference point property.

 (\Leftarrow) Assume *c* satisfies the reference point property. We build the representation recursively. Consider the set *X*. By the reference point property there exists an element a_1 such that for all subsets of X that contain a_1 , condition α holds. Define *Y* = { $x \in X | c({x, a_1} = x)$. For any $x, y \in Y$, $x \neq y$, define $x \succ_{a_1} y$ if $x = c({a_1, x, y})$ (including the case $y = a_1$), and for any $x \notin Y$ define $a_1 \succ_{a_1} x$. Extend \succ_{a_1} such that the elements in $X\ Y$ are ordered arbitrarily. This preference relation, \succ_{a_1} , is well defined by condition α , and it rationalizes the choices of c whenever a_1 is available. Now consider the set $X \setminus \{a_1\}$, and repeat the procedure. We'll find an a_2 and \succ_{a_2} such that ϵ_a represents all choices of *c* whenever a_2 is available in the subsets of $X \setminus \{a_1\}$. For completeness, assume that for any $i \neq 1$, $a_i \succ_{a_2} a_1$. Also construct the *O* − *preference* such that $a_1 \succ o$ a_2 . By induction we can complete the *O* − *preference*, and \succ_a for every *a* in X.

b. Try to come up with a procedure satisfying the reference point axiom which is not stated explicitly in the language of the lexicographical rational choice function (No idea about the answer).

problem 8:

Consider a decision maker who has in mind a set of rationales and a preference relation and chooses the best alternative that he can rationalize.

Formally, we say that a choice function *c* **is rationalized if there is an asymmetric complete relation (not necessarily transitive!) and a set of partial orderings** $\{k_k\}_{k=1...K}$ (called rationales) such that $c(A)$ is the \geq -maximal **alternative from among those alternatives found to be maximal in** *A* **by at least one rationale (given a binary relation** \triangleright we say that x is \triangleright -maximal in A if $x \succ y$ for all $y \in A$). Assume that the relations are such that the procedure **always leads to a solution.**

We say that a choice function *c* **satisfies The Weak Weak Axiom of Revealed Preference (WWARP) if for all** $\{x, y\} \subset B_1 \subset B_2$ ($x \neq y$) and $c\{x, y\} = c(B_2) = x$, then $c(B_1) \neq y$.

a. Show that a choice function satisfies WWARP if and only if it is rationalized. For the proof, construct rationales, one for each choice problem, that are asymmetric binary relations and allow that will not necessarily be transitive.)

 (\Leftarrow) Let us see first that the axiom is satisfied by any rationalized choice function: If x is chosen from B_2 then has a rationale in B_2 (I.e. there is a rationale k_k such that x is the k_k -maximal in B_2). Thus, it has a rationale also in B_1 . If y were chosen from B_1 , then it has a rationale in B_1 as well. Since γ is chosen from B_1 it must be that $y > x$. For $\{x, y\}$ both x and y have rationales and thus y would have been chosen from $\{x, y\}$, a contradiction.

 (\Rightarrow) Let *c* be a choice function satisfying WWARP. For every set *B*, define $x \succ_B y$ iff $x = c(B)$ and $y \in B$. Obviously, this rationale is a very partial ordering. As to the top preferences \rightarrow , they are elicited by the choice from the two-element sets: $x \rightarrow y$ if $C({x,y}) = x$.

To see that those definitions "work", assume $c(B_1) = x$ but there is rationale \succ_{B_2} and a *y* which is \succ_{B_2} -maximal in B_1 such that $y \succ x$. It must be that $B_1 \subset B_2$ and $c(B_2) = y$. By definition of \ge also $C(\{x,y\}) = y$. A contradiction to *WWARP*.

b. What do you think about the axiomatization?

There might be other ways in which people's choices satisfy WWARP. Axiomatizing such a choice with partial orderings for each subset might be representing the choice procedure in a much more complex way than actual.

Consider the "warm-glow" procedure: The decision maker has two complete orderings in mind: one moral \gtrsim_M and one selfish \gtrsim_S . He chooses **the most moral alternative** *m* **as long as he doesn't "lose" too much by not choosing the most selfish alternative. Formally, for every alternative** *s* **there is some alternative** $l(s)$ such that if the most selfish alternative is s then he is **willing to choose** *m* **as long as** $m \geq s$ *l*(*s*). If $l(s) \geq s$ *m*, he chooses *s*.

The function *l* satisfies $s \succsim_{S} l(s)$ and $s \succsim_{S} s'$ iff $l(s) \succsim_{S} l(s')$.

d. Show that WWARP is satisfied by this procedure.

Assume in contradiction that WWARP is violated, i.e. there exists ${x, y}$ ∈ *B*₁ ⊂ *B*₂ (*x* ≠ *y*) and *C*({*x,y*}) = *C*(*B*₂) = *x*, and *C*(*B*₁) = *y*.

Assume x is the moral maximal in B_2 . Clearly x is also the moral maximal in B_1 and this implies that y is the selfish maximal in B_1 . Since x is chosen in B_2 , it must be that $x \succeq s l(s_{B_2})$, where s_{B_2} is the selfish maximal in B_2 . But $B_1 \subset B_2$ implies $s_{B_2} \succeq_s s_{B_1} = y$. By monotonicity of $l(s), x \succeq_s l(y)$ which contradicts that $c(B_1) = y$.

Assume that x is the selfish maximal in B_2 . Clearly x is also the selfish maximal in B_1 , hence *y* is the moral maximal in B_1 . This implies that in $\{x, y\}$, *x* is the selfish maximal and *y* is the moral maximal. Since *x* is chosen in $\{x, y\}$, it must be that $y \not\geq s$ *l(x)*. But since *x* is also selfish maximal in B_1 this contradicts that $c(B_1) = y$.

e. Show directly that the warm-glow procedure is rationalized (in the sense of the definition in this problem).

There are two rationales, the selfish and the moral orderings. The final relation $>$ is the moral ordering, that is $x > y$ if $x >_m y$. However, if $l(y) >_s x$, then it is reversed, that is $y > x$.

To see that this works, given any set we choose the moral and selfish maximal, *m* and *s*. Then we apply the final ordering. Note that if $m \geq s$ *l(s)* then the ordering says $m > s$. Otherwise, $s > m$ as desired.

Problem Set 4 – Consumer Preferences

Problem 1. Consider the preference relations on the interval [0,1] which are continuous. What can you say about those preferences which are also strictly convex?

We will show that a continuous preference relation \succeq on $X = [0,1]$ is strictly convex iff there exists a point x^* such that $b > a$ for all $a < b \le x^*$ or all $x^* \ge b > a$.

(a) Let \succeq be continuous and strictly convex. Since the preferences are continuous and *X* is compact there exists a unique $x^* \in X$ that maximizes the preferences (see Lecture 5). Let $0 \le a < b \le x^*$. By definition $a \precsim x^*$ and $b = \alpha a + (1 - \alpha)x^*$ for some $\alpha \in [0, 1)$ and thus, by strict convexity $a \prec b$. The case, for two points in $[x^*, 1]$ is analogous.

(b) Assuming that the preferences are increasing in $[0, x^*]$ and decreasing in *[x*[∗], 1], we will show that they satisfy strict convexity. Let α ∈ (0, 1) and $a, b \in X$ be such that $a \neq b$ and $a \geq b$. It must be that $\alpha a + (1 - a)b$ is either between a and x^* or between *b* and x^* . If $\alpha a + (1 - \alpha)b$ is between *b* and x^* , then $\alpha a + (1 - \alpha)b > b$. If it is between *a* and *x*^{*}, then $\alpha a + (1 - \alpha)b > a \ge b$.

Problem 2. Show that if the preferences satisfy continuity and monotonicity, then the function $u(x)$ defined by $x \sim (u(x), \ldots, u(x))$ is **continuous.**

Let x be a point in X. By definition $u(x) \geq 0$. We need to show that for any $\epsilon > 0$ there exists δ such that $|u(x) - u(y)| < \epsilon$ for any $y \in B_{\delta}(x)$.

If $u(x) - \epsilon \geq 0$, then by monotonicity, $x \succ (u(x) - \epsilon, \ldots, u(x) - \epsilon)$. By continuity, there exists $\delta_1 > 0$ such that $u(y) > u(x) - \epsilon$ for $y \in B_{\delta_1}(x)$.

If $u(x) - \epsilon < 0$, then for any $\delta_1 > 0$, $u(y) > u(x) - \epsilon$ for $y \in B_{\delta_1}(x)$.

Similarly, by monotonicity, $(u(x) + \epsilon, ..., u(x) + \epsilon) > x$ and thus

 $(u(x) + \epsilon, ..., u(x) + \epsilon) > B_{\delta_2}(x)$ for some $\delta_2 > 0$ by continuity. Therefore, $u(x) + \epsilon > u(y)$ for $y \in B_{\delta_2}(x)$.

Define $\delta = \min\{\delta_1, \delta_2\}$. Then, $|u(x) - u(y)| < \epsilon$ for any $y \in B_\delta(x)$.

Problem 3. In a world with two commodities, consider the following condition: The preference relation \succeq satisfies Convexity 4 if for all x and $\epsilon > 0$

$$
(x_1,x_2) \sim (x_1 - \epsilon, x_2 + \delta_1) \sim (x_1 - 2\epsilon, x_2 + \delta_1 + \delta_2)
$$
 implies $\delta_2 \ge \delta_1$.

Interpret Convexity 4 and show that for strong monotonic and continuous preferences, it is equivalent to the convexity of the preference relation.

Interpretation: If after an x_1 is reduced by ϵ , the consumer must be compensated with δ units of good 2 in order to remain indifferent to x, then he must be compensated with *at least* 2δ units of good 2 if his consumption of x_1 is decreased by 2ϵ .

Convexity 1 Convexity 4: Let

$$
(x_1, x_2) \sim (x_1 - \epsilon, x_2 + \delta_1) \sim (x_1 - 2\epsilon, x_2 + \delta_1 + \delta_2).
$$
 By convexity 1,

$$
(x_1 - \epsilon, x_2 + \frac{\delta_1 + \delta_2}{2}) = \frac{1}{2}(x_1, x_2) + \frac{1}{2}(x_1 - 2\epsilon, x_2 + \delta_1 + \delta_2)
$$

$$
\ge (x_1 - \epsilon, x_2 + \delta_1).
$$

Then, $(\delta_1 + \delta_2)/2 \ge \delta_1$ by monotonicity and thus $\delta_2 \ge \delta_1$.

Convexity 4 \Rightarrow **Convexity 1:** First, we show that if $x \sim y$, then $(x + y)/2 \succeq y$. If $x \neq y$ then by strong monotonicity we can WLOG assume $x_1 > y_1$ and $y_2 > x_2$. Define $\Delta > 0$ by $\Delta = (y_2 - x_2)/2$ and $\epsilon = (x_1 - y_1)/2$. By strong monotonicity

$$
(x_1 - \epsilon, x_2 + 2\Delta) = (\frac{x_1 + y_1}{2}, y_2) > y \sim x
$$

> $(\frac{x_1 + y_1}{2}, x_2) = (x_1 - \epsilon, x_2).$

By continuity, there exists $\delta > 0$ such that

$$
(x_1,x_2) \sim (x_1-\epsilon,x_2+\delta) \sim y = (x_1-2\epsilon,x_2+2\Delta).
$$

By Convexity 4, $2\Delta - \delta \ge \delta$ and thus $\Delta \ge \delta$. By monotonicity,

$$
\frac{x+y}{2}=(x_1-\epsilon,x_2+\Delta)\gtrsim (x_1-\epsilon,x_2+\delta)\sim y.
$$

Now if $x \geq y$, then there exists *z* on the interval which connects 0 and *x*, such that $z_k \le x_k$ for all k and $z \sim y$. Then, by monotonicity and the previous result, $(x + y)/2 \geq (z + y)/2 \geq y$. The rest follows from the following Lemma:

Lemma: If \succeq are continuous preferences, then \succeq are convex iff $[x \succeq y$ implies $(x + y)/2 \ge y$ for all $x, y \in X$.

Proof: Assume $x \geq y$ and $z = \alpha x + (1 - \alpha)y$ for $\alpha \in [0, 1]$. We will show that $z \geq y$. Construct a sequence $\{(x^n, y^n)\}$ such that both $x^n, y^n \geq y$ and *z* between x^n and y^n . Define $x^0 = x$, $y^0 = y$. Continue inductively. Let $m^n = (x^n + y^n)/2$. Then, m^n is at least as good as either x^n or y^n and the above argument and transitivity imply that it is at least as good as *y*. Define:

> $x^{n+1} = m^n$ and $y^{n+1} = y^n$ if *z* lies between y^n and m^n , and $x^{n+1} = x^n$ and $y^{n+1} = m^n$ otherwise.

Thus, x^{n+1} , $y^{n+1} \ge y$ and *z* is between x^{n+1} and y^{n+1} . Since y^n , $x^n \to z$, $z \ge y$ by

continuity.

Problem 4. Complete the proof (for all *K***) of the claim that any continuous preference relation satisfying strong monotonicity quasi-linearity in all** $\boldsymbol{\epsilon}$ ommodities can be represented by a utility function of the form $\sum_{k=1}^K \alpha_k x_k$, **where** $\alpha_k > 0$ for all *k***.**

Proof by induction on *K*: We have already proved this for $K = 1$ and 2.

Let \succsim be a preference relation satisfying the problem's assumptions. Consider the preferences restricted to the set of all vectors of the type $(0, x_2, \ldots, x_K)$. The preferences satisfy Continuity, Strong Monotonicity and Quasi-Linearity in goods 2,..,*K*. By the induction hypothesis, there is a vector of positive numbers $(\alpha_k)_{k=2,\dots,K}$ such that $(0, x_2,..., x_K) \sim (0, \sum_{k=2}^K \alpha_k x_k, 0,..., 0).$

By quasi-linearity in good 1, $(x_1, x_2,..., x_K) \geq (y_2, y_2,..., y_K)$ iff $(x_1, \sum_{k=2}^K \alpha_k x_k, 0, \ldots, 0) \ge (y_2, \sum_{k=2}^K \alpha_k y_k, 0, \ldots, 0).$

The relation over all vectors of the type $(x_1, x_2, 0, \ldots, 0)$ satisfies the three properties in the first two dimensions. Thus, there exists $\beta_1, \beta_2 > 0$ such that $(x_1, x_2, 0, \ldots, 0) \sim (\beta_1 x_1 + \beta_2 x_2, 0, 0, \ldots, 0)$ and thus $x \sim (\beta_1 x_1 + \sum_{k=2}^K \beta_2 \alpha_k x_k, 0, \ldots, 0)$ and by strong monotonicity in the first good, the preferences have a linear utility representation.

Problem 5. Show that for any consumer's preference relation satisfying continuity, monotonicity, strong monotonicity with respect to commodity 1 and quasi-linearity with respect to commodity 1, there exists a number $v(x)$ **such that** $x \sim (v(x), 0, \ldots, 0)$ for every vector x **.**

Since \succsim satisfies continuity and monotonicity every bundle is indifferent to a bundle on the main diagonal. Thus, it is sufficient to show the claim for bundles on the main diagonal.

Let $e = (1, \ldots, 1)$ and define

$$
T = \{ \alpha \in \mathfrak{R}_+ \mid \alpha e \succ (x_1, 0, \ldots, 0) \text{ for all } x_1 \in \mathfrak{R}_+ \}.
$$

We will see that $T = \emptyset$. Assume that $T \neq \emptyset$. Let $\gamma = \inf T$. There are two cases:

Case 1: $\gamma \in T$. Then $\gamma > 0$ and by strict monotonicity of commodity 1,

 $(1 + \gamma, \gamma, \ldots, \gamma)$ > γe . By continuity, there exists $\epsilon > 0$ such that

$$
(1+\gamma,\gamma-\epsilon,\ldots,\gamma-\epsilon) > \gamma e > (x_1,0,\ldots,0)
$$

for all x_1 .

Since $\gamma - \epsilon < \inf T$, there exists an x_1^* such that

 $(x_1^*,0,\ldots,0) \geq (\gamma - \epsilon, \gamma - \epsilon, \ldots, \gamma - \epsilon)$ and by quasi-linearity in commodity 1,

$$
(x_1^* + 1 + \epsilon, 0, \ldots, 0) \ge (1 + \gamma, \gamma - \epsilon, \ldots, \gamma - \epsilon)
$$
, a contradiction.

Case 2: $\gamma \notin T$. Then $(\beta, 0, \ldots, 0) \ge \gamma e$ for some β . By strong monotonicity of commodity 1, $(\beta + 1,0,...,0)$ > γe . By continuity, there is an $\epsilon > 0$ such that $(\beta + 1,0,\ldots,0)$ > $(\gamma + \epsilon)e$, which contradicts $\gamma = \inf T$.

Problem 6. We say that a preference relation satisfies separability if it can be represented by an additive utility function, that is, a function of the type $u(x) = \sum_{k} v_k(x_k)$.

(a) Show that such preferences satisfy condition S: for any subset of commodities *J***, and for any bundles** *a*,*b*, *c*,*d***, we have**

$$
(a_J,c_{-J}) \succsim (b_J,c_{-J}) \Leftrightarrow (a_J,d_{-J}) \succsim (b_J,d_{-J})
$$

where *xJ*, *y*−*^J* **is the vector that takes the components of** *x* **for any** *k* ∈ *J* **and takes the components of** y **for any** $k \notin J$ **.**

$$
(aJ, c-J) \ge (bJ, c-J) \Leftrightarrow \sum_{k \in J} v_k(a_k) + \sum_{i \notin J} v_i(c_i) \ge \sum_{k \in J} v_k(b_k) + \sum_{i \notin J} v_i(c_i)
$$

$$
\Leftrightarrow \sum_{k \in J} v_k(a_k) + \sum_{i \notin J} v_i(d_i) \ge \sum_{k \in J} v_k(b_k) + \sum_{i \notin J} v_i(d_i)
$$

$$
\Leftrightarrow (aJ, d-J) \ge (bJ, d-J).
$$

Graphically, if two bundles lie on the same horizontal line and $(a, c) \ge (b, c)$, then a change of *c* to *d* will preserve the preference relation, that is $(a,d) \geq (b,d)$.

(b) Show that for *K* 2 **such preferences satisfy the** *Hexagon* − *condition***: If** $(a,b) \geq (c,d)$ and $(c,e) \geq (f,b)$ then $(a,e) \geq (f,d)$.

 $v_1(a) + v_2(b) \ge v_1(c) + v_2(d)$ and

 $v_1(c) + v_2(e) \ge v_1(f) + v_2(b)$ implies

 $v_1(a) + v_2(e) \ge v_1(f) + v_2(d).$

(c) Give an example of a continuous preference relation which satisfies condition S and does not satisfy separability.

Consider any preference relation with linear indifference curves as depicted:

Such preferences violate the Hexagon Condition.

Problem 7. (a) Show that the preferences represented by the utility function $\min\{x_1,\ldots,x_K\}$ are not differentiable.

Let $x^* = (a, \ldots, a)$ and let $v(x^*)$ be a candidate set of subjective values. Without loss of generality, let $v_1(x^*) > 0$. Then, $(1,0,0,...,0) \cdot v(x^*) > 0$ but $(a + \epsilon, a, a, \ldots, a) \sim x^*$ for all ϵ , and thus $(+1,0,0,\ldots,0) \notin D(x^*)$, a contradiction.

(b) Check the differentiability of the lexicographic preferences in \mathbb{R}^2 .

Lexicographic preferences are not differentiable. Let $x \in \mathbb{R}^2$ and assume that $v(x)$ is a vector of subjective values. Since $x + \epsilon(0,1) > x$ for all $\epsilon > 0$, then $(0,1)$ is an improving direction and $v_2(x) > 0$. Then, for small $\delta > 0$, $(-\delta, 1) \cdot v(x) > 0$. However, $(-\delta,1)$ is not an improving direction, a contradiction.

(c) Assume that is monotonic, convex and differentiable such that for every *x*, we have (*) $D(x) = \{d \mid (x+d) > x\}$. What can you say about ≥ 2 ?

We will show that the indifference curves are linear.

By differentiability and (*) there exists $v(x)$ such that $d \cdot v(x) > 0$ iff $x + d > x$. Graphically, any point above the dotted line is strictly better than *x*:

We will show that for any $z \in X$ on the dotted line (that is $zv(x) = xv(x)$), we have $x \sim z$.

First let us see that (**) any $z \in X$ on the dotted line satisfies $x \geq z$. If $z \succ x$, then by (*) $(z - x) \in D(x)$ and by differentiability $(z - x) \cdot v(x) > 0$ but as $zv(x) = xv(x)$, a contradiction.

To see that $z \geq x$, by continuity we can assume that z is not a "corner point". It must be that $v(z) = v(x)$, since otherwise there would be a point on $\{y \mid yv(z) = zv(z)\}$ such that $yv(x) > xv(x)$ but by (*) $y > x$ though $x \ge z \ge y$. Thus $(x - z) \cdot v(z) = 0$ and by (**) (applied to *z*) $z \ge x$.

(d) Assume that is a monotonic, convex and differentiable preference relation. Let $E(x) = \{d \in \Re^K \mid \text{there exists } \epsilon^* > 0 \text{ such that } x + \epsilon d \prec x \text{ for all }$ $\epsilon \leq \epsilon^*$. Show that $\{-d \mid d \in D(x)\} \subseteq E(x)$ but not necessarily $\{-d | d \in D(x)\} = E(x)$.

We first show that $\{-d \mid d \in D(x)\} \subseteq E(x)$. By contradiction, let $d \in D(x)$ be such that $-d \notin E(x)$. WLOG $x + d > x$ and $x - d ≥ x$. By definition of $D(x)$, $d \cdot v(x) > 0$ and $e \cdot v(x) > 0$ for some *e* with $e_k \leq d_k$ with at least one strict inequality. For $\epsilon > 0$ small enough $x + \epsilon e > x$. By convexity any convex combination of $x + \epsilon e$ and *x* − *d* is at least as good as *x* but the segment contains points which by monotonicity are at least as bad as *x*.

Let \geq be represented by $u(x) = x_1x_2$. Since *u* is quasi-concave, has continuous partial derivatives and satisfies $u_i(x) > 0$. Thus, the relation \geq is convex, monotonic and differentiable. Let $d = (1,−1)$ and note that $-d \in E(2,2)$ but $d \notin D(2,2)$.

(e) Consider the consumer's preferences in a world with two commodities defined by:

$$
\mathbf{u}(\mathbf{x}_1, \mathbf{x}_2) = \left\{ \begin{array}{ll} x_1 + x_2 & \text{if } x_1 + x_2 \le 1 \\ 1 + 2x_1 + x_2 & \text{if } x_1 + x_2 > 1 \end{array} \right. .
$$

Show that these preferences are not continuous but nevertheless are differentiable according to our definition.

If $x_1 + x_2 \le 1$, then differentiability holds for $v(x) = (1,1)$ and if $x_1 > 1$, then differentiability holds for $v(x) = (2,1)$. The preferences are not continuous, since $(0,2)$ > $(1,0)$, but $(0,2)$ \prec $(1+\epsilon,0)$ for ϵ > 0.
Problem Set 5– Demand: Consumer Choice

Problem 1. Verify that when preferences are continuous, the demand function $x(p, w)$ is continuous in prices and in wealth (and not only in p).

Let $\{(p^n, w^n)\}$ converge to (p, w) . Since $x(p, w)$ is homogeneous of degree zero, then

$$
x(p^n, w^n) = x(\frac{p^n}{w^n}, 1).
$$

Since demand is continuous in *p*, then

$$
x(\frac{p^n}{w^n},1)\to x(\frac{p}{w},1)=x(p,w),
$$

where the equality follows from $x(p, w)$ being homogeneous of degree zero.

Problem 2. Show that if a consumer has a homothetic preference relation, then his demand function is homogeneous of degree one in *w***.**

Let $\lambda > 0$ and $y \in B(p, \lambda w)$. Then $y/\lambda \in B(p, w)$. Since $x(p, w) \ge y/\lambda$ and preferences are homothetic, then $\lambda x(p,w) \geq y$, and thus, $\lambda x(p,w)$ is the best element in *B*(*p*, λw), that is $x(p, \lambda w) = \lambda x(p, w)$.

Problem 3. Consider a consumer in a world with $K = 2$, who has a **preference relation that is convex and quasi-linear in the first commodity. How does the demand for the first commodity change with** *w***?**

Claim: For any *p* either there is no *w* such that $x_1(p, w) > 0$ or there exists an *w*[∗] ≥ 0 such that if *w* ≤ *w*[∗], then the consumer does not consume the first commodity, and if $w \geq w^*$, then the first commodity absorbs all changes in wealth, that is $x_1(p, w) = \frac{w - w^*}{p_1}$.

Proof: Normalize $p_1 = 1$. First, we show if $x_1(p, w) = 0$, then $x_1(p, w') = 0$ for $w' \leq w$.

Let $w' \leq w$. Denote $a = (0, w'/p_2)$, $b \in B(p, w')$ such that $pb = w'$, *c* = $(a_1 + (w - w'), a_2)$, *d* = $(b_1 + (w - w'), b_2)$ and *e* = $(0, w/p_2)$.

Note that $pd = w$ and c is between e and d. Since $e \geq d$ then $c \geq d$. By the quasi-linearity $a \geq b$. Therefore $x(p, w') = a = (0, w'/p_2)$.

Now, if there is *w* such that $x_1(p, w) > 0$ then by continuity of the demand function $w^* = \max\{w \mid x_1(p,w) = 0\}$ exists. We need to show that *x*(*p*,*w*) = (*w* − *w*[∗],*w*^{*}/*p*₂) for all *w* ≥ *w*^{*}. If *x*₂(*p*,*w*) > *w*^{*}/*p*₂, for some *w* > *w*^{*}, let $a = x(p, w)$ and by the quasi linearity in the first commodity $b = (0, a_2) \geq y$ for all $y \in B(p, p_2x_2(p, w))$ contradicting the definition of w^* (see graph). If $x_2(p, w) < w^*/p_2$, for some $w > w^*$, then $c = x(p, w)$ and

 $d = (x_1(p,w) - (w-w^*), x_2(p,w)) \ge e = (0, w^*/p_2)$, a contradiction to the definition of *w*[∗].

Problem 4. Let \succeq be a continuous preference relation (not necessarily strictly convex) and w a number. Consider the set $G = \{(p,x) \in \mathbb{R}^K_{++} \times \mathbb{R}^K_{+} \mid x \text{ is }$ **optimal in** $B(p, w)$. (For some price vectors, there could be more than one $(p, x) \in G$ **.) Calculate** *G* for the case of $K = 2$ and preferences represented by $x_1 + x_2$.

Show that for any preference relation, *G* **is a closed set.**

If preferences are represented by $x_1 + x_2$, then the consumer will buy the cheapest good. Therefore, *G* is the set

 $\{(p, (0, w/p_2)) | p_2 < p_1\} \cup \{(p, (w/p_1, 0)) | p_1 < p_2\} \cup \{(p, (a, w/p_2 - a) | p_2 = p_1, a \in [0, w/p_2]\})$ Let (p^n, x^n) be a sequence of points in *G* converging to (p, x) . Then $p^n x^n \leq w$ for every *n*, and thus $px = \lim_{n\to\infty} p^n x^n \leq w$, i.e. $x \in B(p, w)$.

If *x* is not optimal in *B*(p, w), then there exists a $y \in B(p, w)$ such that $y \succ x$. By continuity, there exists an $\epsilon > 0$ such that $B_{\epsilon}(y) > B_{\epsilon}(x)$, and thus there exists a bundle $z \in B_{\epsilon}(y)$ such that $pz < w$ and $z > x$. Since $p^n \to p$, then $p^n z \leq w$ for *n* large. Since $x^n \to x$, then $x^n \in B_\epsilon(x)$ for *n* large, and thus $z \succ x^n$, a contradiction to x^n be optimal in $B(p^n, w)$.

Problem 5. Determine whether the following consumer behavior patterns are fully rationalized (assume K=2):

a. The consumer's demand function is $x(p, w) = (2w/(2p_1 + p_2), w/(2p_1 + p_2)).$ Yes. $x(p, w)$ is rationalized by the monotonic preferences represented by $u(x) = \min\{x_1, 2x_2\}.$

b. The consumer consumes up to quantity 1 of commodity 1 and spends his excess wealth on commodity 2.

Yes. $x(p, w)$ is rationalized by the monotonic preferences represented by

$$
u(x) = \begin{cases} x_1 & \text{if } x_1 < 1 \\ 1 + x_2 & \text{if } x_1 \ge 1 \end{cases}
$$

c. The consumer chooses the bundle (x_1, x_2) which satisfies $\frac{x_1}{x_2} = \frac{p_1}{p_2}$ and costs w **. (Does the utility function** $u(x) = x_1^2 + x_2^2$ rationalize the consumer's **behavior?)**

No. The behavior violates the WA. $x((2,1),5) = (2,1)$ and $x((1,2),5) = (1,2)$. Both bundles are affordable in both budget sets.

The function $u(x) = x_1^2 + x_2^2$ does not rationalize $x(p, w)$ since a consumer maximizing *u* would allocate all wealth to the cheapest good. The "first order condition" approach is not appropriate because preferences represented by *u* are not convex.

Problem 6. In this question, we consider a consumer who behaves differently from the classic consumer we talked about in the lecture. Once again we consider a world with *K* **commodities. The consumer's choice will be from budget sets. The consumer has in mind a preference relation that satisfies continuity, monotonicity, and strict convexity; for simplicity, assume it is represented by a utility function** *u***.**

The consumer maximizes utility up to utility level $u⁰$. If the budget set **allows him to obtain this level of utility, he chooses the bundle in the budget set with the highest quantity of commodity 1 subject to the constraint that** his utility is at least u^0 .

a. Formulate the consumer's problem.

 $\max_{x \in B(p,w)} u(x)$ if $\max_{x \in B(p,w)} u(x) < u^0$, and $\max_{x \in B(p,w)} x_1$ s.t. $u(x) \ge u^0$ if $\max_{x \in B(p,w)} u(x) \ge u^0$.

b. Show that the consumer's procedure yields a unique bundle.

If $\max_{x \in B(p,w)} u(x) < u^0$, then the consumer acts as in the standard framework. $x(p, w)$ exists because preferences are continuous and is unique because preferences are strictly convex.

If $\max_{x \in B(p,w)} u(x) \geq u^0$, define $\tilde{B} = \{x \in B(p,w) \mid u(x) \geq u^0\}$, which is compact, and convex (by strict convexity). Then $\max_{x \in \tilde{B}} x_1$ exists. If both y and z are solutions then by the strict convexity $u((y+z)/2) > u_0$ and thus there is a vector x such that $u(x) > u^0$ and $x_1 > y_1$ contradicting the optimality of *y* in \tilde{B} .

c. Is this demand procedure rationalizable?

Yes. The procedure is rationalized by

$$
v(x) = \begin{cases} u(x) & \text{if } u(x) < u^0 \\ u^0 + x_1 & \text{if } u(x) \ge u^0. \end{cases}
$$

d. Does the demand function satisfy Walras Law?

Yes. Preferences are monotonic.

e. Show that in the domain of (p, w) for which there is a feasible bundle **yielding utility of at least** *u*⁰ **the consumer's demand function for commodity 1** is decreasing in p_1 and increasing in w .

In both cases the budget set is enlarging and the consumer could obtain more x_1 and preserve u^0 .

f. Is the demand function continuous?

Yes. By question 2, showing that the demand $x(p, w)$ is continuous in p is sufficient. Let $z(p, w)$ be the solution of $\max_{x \in B(p, w)} u(x)$. Note that in this case $z(p, w)$ is not necessarily the consumer's demand $x(p, w)$.

Let $\{p^n\}$ converge to p. If $u(z(p, w)) < u^0$ then for *n* large enough

max_{x∈B} p^n _{*w*} $u(x)$ < u^0 and the demand is $z(p^n, w)$ converges to $z(p, w)$ which is the demand in (p, w) .

Assume $u(z(p, w)) \geq u^0$. Let $m = \inf_{i,n} p_i^n > 0$, the infimum of the commodity prices. Then $x(p^n, w) \in [0, w/m]^K$ for all *n*, and thus WLOG we can assume that $x(p^n, w)$ converges to a bundle *y*. By contradiction, assume that $y \neq x(p, w)$.

If $u(y) < u^0$, then by continuity, there exists an $\epsilon > 0$ such that $B_{\epsilon}(x(p,w)) > B_{\epsilon}(y)$. Then, there exists a $a \in B_{\epsilon}(x(p,w))$ such that $pa < w$ and $a > y$. For *n* large, $p^n a \leq w$ and $a > x(p^n, w)$, a contradiction.

If $u(y) \ge u^0$, then $x_1(p, w) > y_1$. Let $a = \frac{1}{2}x(p, w) + \frac{1}{2}y$. Then $a_1 > y_1$, $a \in B(p, w)$ and $u(a) > u^0$ by strict convexity. By continuity, there exists an $\epsilon > 0$ small such that $a_1 - \epsilon > y_1 + \epsilon$, $p \cdot (a - \epsilon e_1) < w$ and $u(a - \epsilon e_1) \ge u^0$. Thus for *n* large, $B(p_n, w)$ contains $a - \epsilon e_1$ which yields utility larger than u^0 and quantity larger than $x_1(p^n, w)$, a contradiction.

Problem 7. It's a common practice in economics to view aggregate demand as being derived from the behavior of a "representative consumer". Give two examples of "well-behaved" consumer preference relations that can induce average behavior that is not consistent with maximization by a "representative consumer". (That is, construct two "consumers", 1 and 2, who choose the bundles x^1 and x^2 out of the budget set *A* and the bundles y^1 and y^2 out of the budget set B so that the choice of the bundle $\frac{x^1+x^2}{2}$ from A and of the bundle $\frac{y^1+y^2}{2}$ from B is inconsistent with the model of the rational **consumer).**

Let
$$
(p^A, w^A) = ((1, 2), 8), (p^B, w^B) = ((2, 1), 8)
$$
 and
\n
$$
u_1(x) = \begin{cases} x_1 & \text{if } x_1 < 4 \\ 4 + x_2 & \text{if } x_1 \ge 4 \end{cases} \qquad u_2(x) = \begin{cases} x_2 & \text{if } x_2 < 4 \\ 4 + x_1 & \text{if } x_2 \ge 4 \end{cases}
$$

The demands of the two agents in A will be $(4,2)$ and $(0,4)$ and thus

 $\overline{x^A}(p,w) = (2,3)$. Similarly, $\overline{x^B}(p,w) = (3,2)$. Both average bundles are interior in *A* and in *B*. Thus, we the average demand violates the weak axiom.

Problem 8. A commodity *k* **is Giffen if the demand for the** *k***'th good is increasing in** p_k **. A commodity** k **is inferior if the demand for the commodity decreases with wealth. Show that if there is a vector** (p, w) such that the **demand for the** *k***'th commodity is rising after its price has increased, then there is a vector** *p*′ ,*w*′ **such that the demand of the** *k***'th commodity is falling after the income has increased (Giffen implies inferior).**

Let e_k be the vector with the k th component being 1 and all other components being 0. We have $x_k(p + \epsilon e_k, w) > x_k(p, w)$. Let $w' \geq w$ be the "compensating" wealth level, that is $[p + \epsilon e_k] \cdot x(p, w) = w'$. Thus, $x(p + \epsilon e_k, w') \ge x(p, w)$.

By definition

 $px(p + \epsilon e_k, w') + \epsilon x_k(p + \epsilon e_k, w') = [p + \epsilon e_k] \cdot x(p + \epsilon e_k, w') \leq w' = w + \epsilon x_k(p, w).$

If $x_k(p + \epsilon e_k, w') > x_k(p, w)$ then $px(p + \epsilon e_k, w') < w$ contradicting the optimality of $x(p, w)$ in $B(p, w)$.

Problem Set 6 - Choice over Budget Sets and the Dual Consumer

Problem 1. In a world with two commodities, consider a consumer's preferences that are represented by the utility function $u(x_1, x_2) = \min\{x_1, x_2\}$. **a. Calculate the consumer's demand function.**

A solution x^* must satisfy $x_1^* = x_2^*$ and $px^* = w$. Thus, $x(p, w) = (w/(p_1 + p_2), w/(p_1 + p_2)).$

b. Verify that the preferences satisfy convexity.

Let $u(y)$, $u(z) \ge u(x)$, that is $\min\{y_1, y_2\} \ge \min\{x_1, x_2\}$ and $\min\{z_1, z_2\} \geq \min\{x_1, x_2\}$. Then $u(\alpha y + (1 - \alpha)z) = \min{\alpha y_1 + (1 - \alpha)z_1, \alpha y_2 + (1 - \alpha)z_2} \ge$

 $\alpha \min\{y_1, y_2\} + (1 - \alpha) \min\{z_1, z_2\} \ge \min\{x_1, x_2\} = u(x).$

c. Calculate the indirect utility function $v(p, w)$.

 $v(p, w) = u(x(p, w)) = w/(p_1 + p_2).$

d. Verify Roy's Identity.

$$
x_i(p,w) = -\frac{\partial v(p,w)/\partial p_i}{\partial v(p,w)/\partial w} = -\frac{-w/(p_1+p_2)^2}{1/(p_1+p_2)} = \frac{w}{p_1+p_2}.
$$

e. Calculate the expenditure function *ep*,*z* **and verify the Dual Roy's Identity.**

Obviously, $h_i(p, z) = min\{z_1, z_2\}$. Thus, $e(p, z) = min\{z_1, z_2\}(p_1 + p_2)$. Then $h_i(p, z) = \partial e(p, z)/\partial p_i = min\{z_1, z_2\}.$

Problem 2. Imagine that you are reading a paper in which the author uses the indirect utility function $v(p_1, p_2, w) = w/p_1 + w/p_2$. You suspect that the **author's conclusions in the paper are the outcome of the"fact" that the function** *v* **is inconsistent with the model of the rational consumer. Take the following steps to make sure that this is not the case:**

a. Use Roy's Identity to derive the demand function.

$$
x_i(p,w) = -\frac{\partial v(p,w)/\partial p_i}{\partial v(p,w)/\partial w} = -\frac{-w/p_i^2}{(p_1+p_2)/p_1p_2} = \frac{wp_j}{p_i(p_1+p_2)}.
$$

b. Show that if demand is derived from a smooth utility function, then the indifference curve at the point (x_1, x_2) has the slope $-\sqrt{x_2}/\sqrt{x_1}$.

By part (a), $x_i(p, w) > 0$ for $i = 1, 2$. Note that $x_2(p, w) = (\frac{p_1}{p_2})^2 x_1(p, w)$. If *u* is quasi-concave, then

$$
\frac{\partial u(x)/\partial x_1}{\partial u(x)/\partial x_2} = \frac{p_1}{p_2} = \sqrt{\frac{x_2}{x_1}}
$$

c. Construct a utility function with the property that the ratio of the partial derivatives at the bundle (x_1, x_2) **is** $\sqrt{x_2}/\sqrt{x_1}$.

 $u(x) = (\sqrt{x_1} + \sqrt{x_2}).$

d. Calculate the indirect utility function derived from this utility function. Do you arrive at the original $v(p_1, p_2, w)$? If not, can the original indirect utility **function still be derived from another utility function satisfying the property in (c)?**

The indirect utility function derived from *u* is $u(x(p, w)) = (\sqrt{wp_1/p_2(p_1 + p_2)} + \sqrt{wp_2/p_1(p_1 + p_2)}) = \sqrt{w(p_1 + p_2)} / \sqrt{p_1 p_2}$. The function $u^2(x)$ represents the same preference relation and $u^2(x(p, w)) = v(p, w).$

Problem 3. A consumer with wealth *w* **is interested in purchasing only one unit of one of the items included in a (finite) set** *A***. All items are indivisible. The consumer does not derive any "utility" from leftover wealth.** The consumer evaluates commodity $x \in A$ by the number V_x (where the value of not purchasing any of the goods is 0). The price of commodity $x \in A$ is $p_x > 0$.

a. Formulate the consumer problem.

Let "*n*" denote not purchasing anything, where $p_n = 0$ and $V_n = 0$. Define *B*(*p*,*w*) = { $x \in A \cup \{n\}$ | $p_x \leq w$ }. We get: $v(p, w) = \max_{x \in B(p, w)} V_x$.

b. Check the properties of the indirect preferences (homogeneity of degree zero, monotonicity, continuity and quasi-convexity).

The proofs of Homogeneity, Monotonicity and Quasi Convexity are valid. The indirect utility function is not continuous: **Take** $K = 1, A = \{x\}$, where $V_x = 1$, $(p_x) = (1)$ and $w = 1$. Then $v(1,1) = 1$ but $v(1,1 - \epsilon) = 0$ for any $\epsilon > 0$.

c. Calculate an indirect utility function for the case in which $A = \{a, b\}$ **and** $V_a > V_b > 0.$

$$
V(p, w) = \begin{cases} V_a & p_a \leq w \\ V_b & p_b \leq w < p_a \\ 0 & p_a, p_b < w \end{cases}
$$

Problem 4. Assume that the consumer's preferences satisfy monotonicity, continuity and strict convexity. Show that the bundle *x* **is the best element in** $B(p, w)$ if and only if for all $y \in B(p, w)$ different then *x* there exists a budget set $B(p',w')$ containing y and not x such that $(p,w) \succ^* (p',w').$

 \Rightarrow Let *x*[∗] = *x*(*p*,*w*). Let *y* ∈ *B*(*p*,*w*), *y* ≠ *x*[∗]. Choose (*p'*,*w'*) such that {*x*|*xp'* = *w'*} is the hyperplane tangent to the indifference curve through *y*. Clearly, $y \in B(p',w')$, *x*[∗] ∉ *B*(p' , w'), and *x*(p , w) ≻ *x*(p' , w').

 \leftarrow Let x^* ∈ $B(p, w)$ such that for any $y \in B(p, w)$ different then x^* there exists a budget set $B(p', w')$ containing y and not x^* such that $(p, w) \succ^* (p', w').$ Then, $x(p,w) > x(p',w') \succsim y$ for all $y \neq x^*$ and thus, it must be that $x(p,w) = x^*$

Problem 5. Show that if the preferences are monotonic, continuous and strictly convex, then the Hicksian demand function $h(p, z)$ is continuous.

Let $\{(p^n, z^n)\}$ converge to (p, z^0) . Define $\bar{z}_k = \sup\{z_k^n\}$, $m = \inf\{p_k^n\}$ and $M = \sup \{ p_k^n \}$. The consumer does not need more than $M\sum$ \bar{z}_k to obtain any z^n .

k $\mathsf{Thus},\, h_l(p^n,z^n)\leq M\sum$ *k* \bar{z}_k/m for all *l*. Thus, WLOG we can assume that $h(p^n,z^n)$ converges to some bundle *h*[∗].

By contradiction, assume that $h^* \neq h(p, z^0)$. By the continuity of the preferences *h*[∗] $\ge z^0$, and thus (assuming that *h*(*p*,*z*) is uniquely defined) *ph*[∗] > *ph*(*p*,*z*⁰). There exists an $\epsilon > 0$ such that $ph^* > p[h(p, z^0) + (\epsilon, \dots, \epsilon)]$ and by monotonicity of the preferences $h(p, z^0) + (\epsilon, \ldots, \epsilon) > z^0$. Then, for *n* large enough, $pⁿh(pⁿ,zⁿ) > pⁿ[h(p,z⁰) + \epsilon e]$ and $h(p,z⁰) + \epsilon e > zⁿ$, a contradiction.

Problem 6. One way to compare budget sets is by using the indirect preferences which involves comparing $\mathbf{x}(p,w)$ and $x(p',w)$. Following are two **other approaches to making such a comparison.**

Define:

$$
CV(p, p', w) = w - e(p', z) = e(p, z) - e(p', z)
$$

where $z = x(p, w)$. This is the answer to the question: What is the change in wealth that would be equivalent, from the perspective of (p, w) , to the change **in price vector from** p **to** p' **?**

Define:

$$
EV(p, p', w) = e(p, z') - w = e(p, z') - e(p', z')
$$

where $z' = x(p',w)$. This is the answer to the question: What is the change in **wealth that would be equivalent, from the perspective of** *p*′ ,*w***, to the change in price vector from** p **to** p' **?**

Now, answer the following questions regarding a consumer in a two-commodity world with a utility function *u***:**

a. For the case of the preferences represented by $u(x_1, x_2) = x_1 + x_2$ **, calculate the two consumer surplus measures.**

$$
CV(p, p', w) = w - \frac{w \min \{p'_1, p'_2\}}{\min \{p_1, p_2\}} = w \frac{\min \{p_1, p_2\} - \min \{p'_1, p'_2\}}{\min \{p_1, p_2\}}
$$

$$
EV(p, p', w) = \frac{w \min \{p_1, p_2\}}{\min \{p'_1, p'_2\}} - w = w \frac{\min \{p_1, p_2\} - \min \{p'_1, p'_2\}}{\min \{p'_1, p'_2\}}
$$

Assume that the price of the second commodity is fixed and that the price vectors differ only in the price of the first commodity.

b. Assume that the first good is a normal good (the demand is increasing with wealth). What is the relation of the two measures to the "area below the demand function" (which is a standard third definition of consumer surplus)?

(This part should be polished)

Let $a = p''_1 < p'_1 = b$ and let A denote the area under the demand curve for commodity 1 between *a* and *b*.

Let $u' = v((a, p_2), w)$ and $u' = v((b, p_2), w)$.

Let $h_1(p, u)$ denote the hicksian demand $h_1(p, z)$ when $u(z) = u$.

If commodity 1 is a normal good, then $h_1(p, u)$ is increasing in u , and thus

$$
h_1((t,p_2),u')\leq h_1((t,p_2),v((t,p_2),w))\leq h_1((t,p_2),u'')\text{ for }t\in[a,b].
$$

Recall that by the Dual Roy's identity, $\frac{\partial e((t,p_2),u)}{\partial p_1} = h_1((t,p_2),u)$ By integrating,

$$
CV(p', p'', w) = \int_a^b \frac{\partial e((t, p_2), u')}{\partial p_1} dt = \int_a^b h_1((t, p_2), u') dt
$$

\n
$$
\leq \int_a^b h_1((t, p_2), v((t, p_2), w)) dt = \int_a^b x_1((t, p_2), w)) dt = A
$$

\n
$$
\leq \int_a^b h_1((t, p_2), u'') dt = \int_a^b \frac{\partial e((t, p_2), u'')}{\partial p_1} dt = EV(p', p'', w).
$$

c. Explain why the two measures are identical if the individual has quasi-linear preferences in the second commodity and in a domain where the two commodities are consumed in positive quantities.

Recall that by Question 4 of PS5, the consumer's demand adjusts to a change in wealth by adjusting consumption of the quasi-linear good (in this case commodity 2) and the consumption of commodity 1 is constant. Consider the following diagram where CV and EV are drawn. Since the indifference curves are parallel horizontally, the two measures are identical.

Problem 7.

a) Verify that you know the envelope theorem, which states conditions under which the following is correct:

Consider a maximization problem $\max_x \{u(x, \alpha_1, \ldots, \alpha_n) \mid g(x, \alpha_1, \ldots, \alpha_n) = 0\}.$ Let *V*($\alpha_1, \ldots, \alpha_n$) be the value of the maximization. Then $\frac{\partial V}{\partial \alpha_i}(a_1, \ldots, a_n) =$ $\frac{\partial (u-\lambda g)}{\partial \alpha_i}(x^*(a_1,,\dots,a_n),a_1,,\dots,a_n)$ where $x^*(a_1,,\dots,a_n)$ is the solution to the **maximization problem and is the Lagrange multiplier associated with the solution of the maximization problem.**

b) Derive the Roy's identity from the envelope theorem (hint: show that in ${\bf this\ context}\ \frac{\partial V/\partial a_i}{\partial V/\partial a_j}(a_1,,\dots,a_n)=\frac{\partial g/\partial a_i}{\partial g/\partial a_j}(x^*(a_1,,\dots,a_n),a_1,,\dots,a_n)).$

In the context of the consumer's problem we have $(a_1, \ldots, a_n) = (p_1, \ldots, p_k, w)$, $u(x, \alpha_1, \ldots, \alpha_n) = u(x)$ and $g(x, \alpha_1, \ldots, \alpha_n) = w - px$. Note that *u* depends on α_i only through *x* (That is, the prices and wealth don't have a direct effect on the utility. The utility depends on the prices and wealth only through their effect on the bundle *x*). Using the chain rule it implies that $\partial u/\partial \alpha_i(x(\alpha_1,\ldots,\alpha_n),\alpha_1,\ldots,\alpha_n) = \partial u/\partial x \cdot \partial x/\partial \alpha_i$. Thus, if $x^*(a_1,...,a_n)$ is the solution to the maximization problem then $\partial u/\partial x = 0$ and therefore $\partial u / \partial \alpha_i(x^*(\alpha_1, \ldots, \alpha_n), \alpha_1, \ldots, \alpha_n) = 0$ for all *i*.

Denote $\alpha = (a_1, \ldots, a_n)$.

The envelope theorem states that:

$$
\frac{\partial V}{\partial \alpha_i(\alpha)} = \frac{\partial u}{\partial \alpha_i(x^*(\alpha), \alpha)} - \lambda(\alpha) \frac{\partial g}{\partial \alpha_i(x^*(\alpha), \alpha)}
$$

Thus,

$$
\frac{\partial V}{\partial \alpha_i(\alpha)} = -\lambda(\alpha) \frac{\partial g}{\partial \alpha_i(x^*(\alpha), \alpha)}.
$$

By taking ratios and canceling out λ :

$$
\frac{\partial V/\partial \alpha_i}{\partial V/\partial \alpha_j}(\alpha) = \frac{\partial g/\partial \alpha_i}{\partial g/\partial \alpha_j}(x^*(\alpha), \alpha).
$$

Since $g(x, a) = w - px$, for any $i = 1, 2, ..., n - 1$:

$$
\partial g/\partial \alpha_i = \partial g/\partial p_i(x^*(a),a) = -x_i^*
$$

and for $i = n$:

 $\partial g / \partial a_n = \partial g / \partial w (x^*(a), a) = 1$

Plugging this in $\frac{\partial V/\partial \alpha_i}{\partial V/\partial \alpha_j}(\alpha) = \frac{\partial g/\partial \alpha_i}{\partial g/\partial \alpha_j}(x^*(\alpha), \alpha)$ we get $\frac{-\partial v/\partial p_i}{\partial v/\partial w}(\alpha) = x_i^*$ which is exactly Roy's identity.

c) What makes it is easy to prove Roy's identity without using the envelope theorem?

The fact that the utility does not depend directly on the prices (only through the bundle *x*).

Problem Set 7: The Producer

Problem 1. Assume that technology *Z* **and the production function** *f* **describe the same producer who produces commodity** *K* **using inputs**

1,…,*K* − 1**. Show that** *Z* **is a convex set if and only if** *f* **is a concave function.** *Z* is convex

iff $(-v, y), (-v', y') \in Z$, $λ ∈ [0, 1]$ implies that $(-λv – (1 – λ)v', λy + (1 – λ)y') ∈ Z$ iff $y \le f(v)$, $y' \le f(v')$, $\lambda \in [0,1]$ implies that $\lambda y + (1 - \lambda)y' \le f(\lambda v + (1 - \lambda)v')$ iff $\lambda f(v) + (1 - \lambda)f(v') \leq f(\lambda v + (1 - \lambda)v')$ iff *f* is concave.

Problem 2. Calculate the supply function $z(p)$ for each of the following **production functions:**

Let *w* **denote the input prices and** *p* **denote the output price.**

(a) $f(a) = a_1^{\alpha}$, for $\alpha \le 1$

If $\alpha = 1$, then the firm solves $\max_{a_i \geq 0} (p - w)a_1$, and thus $z_1(p, w) = 0$ if $p < w$ and is not defined otherwise.

If $\alpha \in (0,1)$, the firm solves $\max_{a_1 \geq 0} pa_1^{\alpha} - wa_1$, which is strictly concave and strictly increasing at $a_1 = 0$. Then the first order condition is sufficient for maximization, and thus $z_1(p,w) = \left(\frac{w}{ap}\right)^{\frac{1}{a-1}}$

(b) $g(a) = min\{a_1, a_2\}$ $z_1(p,w) = z_2(p,w) = 0$ if $p < w_1 + w_2$ and is not defined otherwise.

Problem 3. Consider a producer who uses *L* **inputs to produce** *K* − *L* **outputs. Denote by** *w* the price vector of the *L* inputs. Let $a_k(w, y)$ be the **demand for the** *k***'th input when the price vector is** *w* **and the output vector he wishes to produce is** *y***. Show the following:**

a. $C(\lambda w, y) = \lambda C(w, y)$. $C(\lambda w, y) = \min_{\{a \mid (-a, y) \in Z\}} \lambda wa = \lambda \min_{\{a \mid (-a, y) \in Z\}} wa = \lambda C(w, y)$.

b. *C* is nondecreasing in any input price w_k .

Assume $w'_l \geq w_l$ for all *l*. $C(w', y) = w' a(w', y) \geq wa(w', y) \geq wa(w, y) = C(w, y).$ c. *C* **is concave in** *w***.** Let *w*, *w'* be input prices, $w'' = \lambda w + (1 - \lambda)w'$ for $\lambda \in [0, 1]$. Then $C(w'', y) = [\lambda w + (1 - \lambda)w']a(w'', y) = \lambda wa(w'', y) + (1 - \lambda)w'a(w'', y)$ $\geq \lambda C(w, y) + (1 - \lambda)C(w', y).$

d. **Shepherd's lemma: If** C is differentiable, $\frac{\partial C}{\partial w_k}(w^*,y) = a_k(w^*,y)$ (the *k*th **input commodity).**

Fix *y*. *C* is now a function of *w*. For every *w*, $C(w, y) \leq wa(w^*, y)$. $C(w^*, y) = w^* a(w^*, y)$. Thus $\{(w, c) | c = wa(w^*, y)\}$ is tangent to the graph of the function $C(w, y)$ at $(w^*, C(w^*, y))$. Since *C* is differentiable $\nabla C(w^*, y) = a(w^*, y)$.

e. **If** *C* **is twice continuously differentiable, then for any two commodities** *j* **and** k , $\partial a_j/\partial w_k(w, y) = \partial a_k/\partial w_j(w, y)$.

By Shapherd's Lemma and Young's Theorem (mixed partial derivatives are equal):

$$
\frac{\partial a_j(w,y)}{\partial w_k} = \frac{\partial^2 C(w,y)}{\partial w_k \partial w_j} = \frac{\partial^2 C(w,y)}{\partial w_j \partial w_k} = \frac{\partial a_k(w,y)}{\partial w_j}.
$$

Problem 4. It is usually assumed that the cost function *C* **is convex in the output vector. Much of the research on production has been aimed at investigating conditions under which convexity is induced from more primitive assumptions about the production process. Convexity often fails when the product is related to the gathering of information or data processing.**

Consider, for example, a firm conducting a telephone survey immediately following a TV program. Its goal is to collect information about as many viewers as possible within 4 units of time. The wage paid to each worker is *w* **(even when he is idle). In one unit of time, a worker can talk to one respondent or be involved in the transfer of information to or from exactly one colleague. At the end of the 4 units of time, the collected information must be in the hands of one colleague (who will announce the results).**

Define the firm's product, calculate the cost function and examine its convexity.

The firm's product is units of information.

Denote the agents by 1,...,n. Let $i \rightarrow j$ stands for *i* transfers information to *j* and let → *j* stands for *j* collects information from a viewer. Denote a procedure by a sequence of square brackets, each stands for one period and contains the transfers of information during that period. Agent 1 will be the agent who announces the result.

 $n = 1$. Since there is only one agent, he can use the four units of time only for collecting information from four viewers. Thus, the maximum number of responses is 4. An optimal procedure:

 $\left[\rightarrow 1 \right] \rightarrow 1 \left[\rightarrow 1 \right] \rightarrow 1$

2. The last unit of time has to be used to transfer information from some agent, 2, to agent 1. Therefore, both agents can collect information only for the first three units of time. Thus, the maximum number of responses is 6.

An optimal procedure:

 $\left[\rightarrow 1\rightarrow 2\right] \rightarrow 1\rightarrow 2\left[\rightarrow 1\rightarrow 2\right] \left[\rightarrow 1\right]$

3. Again, the last unit of time has to be used to transfer information from one agent, let us say 2, to agent 1. However, one of these two agents has to get the information from agent 3 one period earlier. Thus, there are two agents who are free to collect information for two periods and one agent who is free for three periods. The total number of collected responses is thus, bounded by 7.

An optimal procedure:

 $[\rightarrow 1, \rightarrow 2, \rightarrow 3] [\rightarrow 1, \rightarrow 2, \rightarrow 3] [\rightarrow 1, 3 \rightarrow 2] [2 \rightarrow 1]$

4. If the firm employs 4 or more agents it can collect at most 8 responses.

Again, the last unit of time has to be used to transfer information from one agent, let us say 2, to 1. It is sufficient to show that each of them cannot hold more

than 4 units of information after three periods. To see it note that after two periods each agent can hold not more than 2 units and thus, after three periods he will have 3 units if he makes a call himself, or 4 units if he gets the information collected earlier by another agent.

An optimal procedure:

 $\left[\rightarrow 1, \rightarrow 2, \rightarrow 3, \rightarrow 4 \right] \left[\rightarrow 1, \rightarrow 2, \rightarrow 3, \rightarrow 4 \right] \left[3 \rightarrow 1, 4 \rightarrow 2 \right] \left[2 \rightarrow 1 \right]$

Let *y* be the "output", the number of responses the center collected and $C(y, w)$ be the minimum cost of producing *y*. Then,

Obviously, *C* is not convex: $C(2, w) = w > w/2 = 5C(0, w) + 5C(4, w)$.

Problem 5. Consider a firm producing one commodity using *L* **inputs, which maximizes production subject to the constraint of nonnegative profits. Show that under reasonable assumptions, the firm's supply function satisfies Homogeneity of degree 0, monotonicity in prices and continuity.**

Let $D(p, w) = \{x \in \Re^L_+ \mid pf(x) - wx \ge 0\}$. The firm solves $\max_{x \in D(p, w)} f(x)$. Let $x(p, w)$ be the firm's input demand and $y(p, w) = f(x(p, w))$ be the firm's optimal output.

Homogeneity of degree zero for $x(p, w)$, $y(p, w)$: It follows from the fact that $D(p, w) = D(\lambda p, \lambda w)$ for $\lambda > 0$.

Optimal output yp, w is non-increasing in w and strictly increasing in p if *f* **is strictly increasing:** If $w' \geq w$, then $D(p, w') \subseteq D(p, w)$, and thus $y(p,w') \leq y(p,w)$. If $p' > p$, then there exists a $x' > x(p,w)$ such that $x' \in D(p',w)$. By *f* strictly increasing, $y(p', w) \ge f(x') > y(p, w)$.

If **f** is continuous and strictly concave, then $x(p, w)$ and $y(p, w)$ are **continuous:** Let $\{(p^n, w^n)\}$ converge to (p, w) . Define $\bar{p} = \sup\{p^n\}$ and $\underline{w} = \inf\{w_k^n\}$, and thus $D(p^n, w^n) \subseteq D(\bar{p}, (\underline{w}, \dots, \underline{w}))$ for all *n*. Then

$$
x^n \in \{x \in \Re^L_+ \mid \bar{p}y(\bar{p}, (\underline{w}, \dots, \underline{w})) - (\underline{w}, \dots, \underline{w}) \cdot x \geq 0\}
$$

since $\bar{p}y(\bar{p}, (w, \ldots, w)) \geq p^n f(x^n)$ and $(w, \ldots, w) \leq w^n$ for all *n*. If $\{x^n\}$ does not converge to $x(p, w)$ then there is a subsequence that converges to some $x^* \neq x(p,w)$. Since *f* is continuous, $pf(x^*) \geq wx^*$. Thus, by strict concavity, *f*(*x*^{*}) < *f*(*x*(*p*,*w*)).

Now (i) $f(x^* + x(p, w)/2) > f(x^*)$ and (ii) $pf(x^*) \ge wx^*$ and $pf(x(p, w)) \ge wx(p, w)$ and by the strict concavity of *f*,

 $pf((x^* + x(p, w))/2) > (pf(x^*) + pf(x(p, w)))/2 \geq w((x^* + x(p, w))/2).$

Therefore we found x' such that $f(x') > f(x^*)$ and $pf(x') - wx' > 0$. Therefore, for *n* large enough, $p^n f(x') - w^n x' \ge 0$ and $f(x') > f(x^n)$, a contradiction.

Problem 6. An event that could have occurred with probability 0.5 **either did or did not occur. A firm must provide a report in the form of "the event occurred" or "the event did not occur". The quality of the report (the firm's product), denoted by** *q***, is the probability that the report is correct. Each of** *k* **experts (input) prepares an independent recommendation which is correct** with probability $1 > p > 0.5$. The firm bases its report on the k **recommendations in order to maximize** *q***.**

a. **Calculate the production function** $q = f(k)$ for (at least) $k = 1, 2, 3, \ldots$

b. We say that a "discrete" production function is concave if the sequence of marginal product is nonincreasing. Is the firm's production function concave?

No. Marginal product is positive from 0 to 1, zero from 1 to 2 and then positive from 2 to 3.

Assume that the firm will get a prize of *M* **if its report is actually correct. Assume that the wage of each worker is** *w***.**

c. Explain why it is true that if *f* **is concave, the firm chooses** *k*[∗] **so that the** *k*[∗]**th worker is the last one for whom marginal revenue exceeds the cost of a single worker.**

The form's profits if it employs *k* workers are: $Mf(k) - kw$. If *f* is concave, then for any worker $k < k^*$, the firm's marginal profit $M[f(k+1) - f(k)] - w$ is positive, whereas for any worker $k > k^*$, the firm's marginal profit is negative.

d. Is this conclusion true in our case?

No. Since the marginal revenue of the second expert is 0, while it is possible that it is optimal for the firm to hire 3 experts.

Problem 7. **Come up with a theory for a producer who maximizes production given the constraint of achieving a level of profit (and does not produce at all if he cannot).**

a. Show conditions under which the producer's problem has a unique solution for every price vector.

Consider there to be only one output, and k inputs $y = f(a_1, \ldots, a_k)$. For a given vector of inputs, and price vector the profit function is $\pi(a,p) = p_y y - p_a a$.

Now we can write the Producer's problem as $max_a f(a)$ s.t. $\pi(a,p) > \rho$. Given prices, if there is no *a* s.t. $\pi(a,p) \geq \rho$, then the producer chooses not to produce.

If the production technology is strictly convex $(f(a))$ strictly concave, decreasing returns to scale), bounded from above, and all prices are strictly positive then we have a unique solution to our problem.

Assume for contradiction that there are two solutions, *a* and *a*′ . It must be that $f(a) = f(a')$ and $\pi(a,p) = \pi(a',p) = \rho$, otherwise we would be able to increase production. Now look at a convex combination of these two points. For $\lambda \in (0,1)$, due to strict concavity of $f(a)$, $f(\lambda a + (1 - \lambda)a') > f(a) = f(a')$, and therefore $\pi(\lambda a + (1 - \lambda)a', p) > \rho$ (since $p_a a = p_a a' = p_a(\lambda a + (1 - \lambda)a')$ and $p_yf(a) = p_yf(a') < p_yf((\lambda a + (1 - \lambda)a'))$. But this means $\lambda a + (1 - \lambda)a'$ is better than *a* and *a*′ .

b. How does the supply of the good change with its price and the price of any of the input goods.

Since we proved above that the solution is unique we can refer to the optimal choice of inputs given prices as $a(p)$.

Consider increasing the output price: let p and p' be price vectors s.t. $p'_y > p_y$ and $p'_a = p_a$. For all a if $p_yf(a) - p_a a \ge \rho$ then $p'_yf(a) - p'_a a > \rho$ and for $\epsilon > 0$ small enough also $p'_{y}f(a+(\epsilon, \ldots, \epsilon))$ $-p'_{a}(a+(\epsilon, \ldots, \epsilon))$ $> \rho.$ Therefore,

 $f(a(p', \rho)) \geq f(a(p, \rho) + (\varepsilon, \ldots, \varepsilon)) \Rightarrow f(a(p, \rho))$ (assuming that *f* is increasing).

Now consider increasing the input price: let p and p' be price vectors s.t. $p'_y = p_y$ and $p'_a > p_a$. Similarly, For all a if $p'_x f(a) - p'_a a \ge \rho$ then $p_x f(a) - p_a a > \rho$ and for $\epsilon > 0$ small enough also $p_y f(a + (\epsilon, \ldots, \epsilon)) - p_a(a + (\epsilon, \ldots, \epsilon)) > \rho$. Therefore, $f(a(p, \rho)) \geq f(a(p', \rho) + (\varepsilon, \ldots, \varepsilon)) \Rightarrow f(a(p', \rho))$ (assuming that *f* is increasing).

c. what can you say about the change in the producer behavior as increases (in the range where he does indeed produce)?

Intuitively, as ρ increases, keeping the prices the same, we shift the price line with same slope up. Due to convexity of the production technology this will lead to less production.

Formally, assume $\rho' > \rho$. For all *a* if $p_yf(a) - p_a a \ge \rho'$ then $p_yf(a) - p_a a > \rho$ and for $\epsilon > 0$ small enough also $p_yf(a+(\epsilon, \ldots, \epsilon)) - p_a(a+(\epsilon, \ldots, \epsilon)) > \rho$. Therefore, $f(a(p, \rho)) \geq f(a(p, \rho') + (\varepsilon, ..., \varepsilon)) > f(a(p, \rho')).$

d. Any other observations you can think of (and prove...)?

Problem Set 8 - Expected Utility

Problem 1. Consider the following preference relations that were described in the text: "the size of the support" and "comparing the most likely prize".

a. Check carefully whether they satisfy axioms *I* **and** *C***.**

Both preference relations violate both axioms:

the size of the support

Let p_t be the lottery: $p_t(z_1) = t$ and $p_t(z_2) = 1 - t$. not *I*: $[z_1] > p_{1/2}$, but for $1 > \alpha > 0$, $\alpha[z_1] \oplus (1 - \alpha)p_{1/2} \sim p_{1/2}$. not *C*: For any $1/n > 0$, $p_{1/n} \sim p_{1/2}$, but in the limit $[z_1] > p_{1/2}$.

comparing the most likely prize

Assume that z_1 is better than z_2 and that "ties are broken in favor" of z_1 : not *I*: $[z_1] \succ [z_2]$, but $p_{\frac{1}{4}} \sim [z_2]$.

not *C*: $(0.5 - 1/n, 0.5 + 1/n) \sim [z_2]$ for all *n* but in the limit $p_{1/2} > [z_2]$.

b. These preference relations are not immune to a certain "framing problem". Explain.

Both preference relations strictly prefer the lottery \$100 with probability 0.4 and \$50 with probability 0.6 to the lottery \$100 with probability 0.4, one blue note of \$50 with probability 0.3 and one green note of \$50 with probability 0.3, even though the lotteries seem to be the "same".

Problem 2. One way to construct preferences over lotteries with monetary prizes is by evaluating each lottery *L* **on the basis of two numbers:** *ExL***, the expectation of** *L***, and** *varL***,** *L***'s variance. Such a construction may or may not be consistent with vNM assumptions.**

a. Show that $u(L) = Ex(L) - (1/4)var(L)$ induces a preference relation that is **not consistent with the vNM assumptions.**

 $[1] \sim 0.5[0] \oplus 0.5[4]$ since $u([1]) = u(0.5[0] \oplus 0.5[4]) = 1$.

However, for $\alpha = 1/2$:

 $\alpha[1] + (1 - \alpha)[0.5[0] \oplus 0.5[2]] > \alpha[0.5[0] \oplus 0.5[4]] + (1 - \alpha)[0.5[0] \oplus 0.5[2]]$ since the utility of the left lottery, 7/8, is greater than the utility of the right lottery, 13/16.

b. Show that $u(L) = Ex(L) - (Ex(L))^2 - var(L)$ is consistent with vNM **assumptions.**

Using the formula $var(X) = Ex(X^2) - (ExX)^2$ we get $u(L) = Ex(L) - (Ex(L))^2 - var(L) =$ $Ex(L) - (Ex(L))^2 - (\sum_{z \in Z} L(z)z^2 - (Ex(L))^2) = \sum_{z \in Z} L(z)(z - z^2)$

is an expected utility function with vNM value $v(z) = z - z^2$.

Problem 3. A decision maker has a preference relation over the space of lotteries *LZ* **having a set of prizes** *Z***. On Sunday he learns that on Monday he will be told whether he has to choose between** *L*¹ **and** *L*² **(probability** $1 > \alpha > 0$) or between L_3 and L_4 (probability $1 - \alpha$). He will make his choice at **that time. Let us compare between two possible approaches the decision maker can take.**

Approach 1: He delays his decision to Monday ("why bother with the decision now when I can make up my mind tomorrow.").

Approach 2: He makes a contingent decision on Sunday regarding what he will do on Monday, that is, he decides what to do if he faces the choice between L_1 and L_2 and what to do if he faces the choice between L_3 and L_4 **("On Monday morning I will be so busy. . .").**

a. Formulate Approach 2 as a choice between lotteries.

The DM chooses one of the four "compound" lotteries in the set

$$
\{\alpha L_i \oplus (1-\alpha)L_j \mid i \in \{1,2\}, j \in \{3,4\}\}.
$$

b. Show that if the preferences of the decision maker satisfy the independence axiom, then his choice under Approach 2 will always be the same as under Approach 1.

Let L_i (L_j) be the preferred lottery in $\{L_1, L_2\}$ ($\{L_3, L_4\}$), and L_{-i} (L_{-j}) be the other lottery. Under approach 1, the DM selects *Li* (*Lj*) if the choice set on Monday is $\{L_1, L_2\}$ ($\{L_3, L_4\}$). Let \succeq be the DM's preferences over the compound lotteries in (a). By *I*,

 $\alpha L_i \oplus (1 - \alpha) L_j \succ \alpha L_i \oplus (1 - \alpha) L_{-i} \succ \alpha L_{-i} \oplus (1 - \alpha) L_{-i}$ and $\alpha L_i \oplus (1 - \alpha) L_j \succ \alpha L_{-i} \oplus (1 - \alpha) L_j$. Thus $\alpha L_i \oplus (1 - \alpha) L_j$ is the best of the "compound" lotteries.

Problem 4. A decision maker is to choose an action from a set *A***. The set of consequences is** *Z***. For every action** *a* ∈ *A***, the consequence** *z*[∗] **is realized** with probability α and any $z \in Z \setminus \{z^*\}$ is realized with probability $r(a,z) = (1 - \alpha)q(a,z)$.

a. Assume that after making his choice he is told that *z*[∗] **will not occur and is given a chance to change his decision. Show that if the decision maker obeys the Bayesian updating rule and follows vNM axioms, he will not change his decision.**

By the vNM Theorem, preferences exhibit expected utility representation. Before learning the information, the DM solves

$$
\max_{a\in A}\left[\sum_{z\in Z\setminus\{z^*\}}r(a,z)v(z)+\alpha v(z^*)\right].
$$

After learning that *z*[∗] will not occur, the DM updates his beliefs so that $r'(a,z) = r(a,z)/(1-\alpha) = q(a,z)$ for $z \in Z \setminus \{z^*\}$ and the DM solves $\max_{a \in A} \sum_{z \in Z \smallsetminus \{z^*\}} r'(a,z) v(z),$ which yields the same solution.

b. Give an example where a decision maker who follows nonexpected utility preference relation or obyes a non-Bayesean updating rule is not time consistent.

Example 1. Assume the DM has a "worst case" preference relation, where z_1 is the best prize, z_2 is the second best and z^* is the worst. Let action a_1 yield z_1 for sure and action *a*² yield *z*¹ and *z*² with equal probability, conditional on *z*[∗] not occurring. Then the DM will initially be indifferent between a_1 and a_2 , but will strictly prefer a_1 after the information is revealed.

Example 2. Assume that $Z = \{1,2,3,z^* = 0\}$ and that $v(z) = z$. Assume that initially his beliefs are: $q(a_1, 2) = 1$, $q(a_2, 3) = 0.4$ and $q(a_2, 1) = 0.6$. Contingentally the DM chooses a_1 . If he updates his beliefs and after he was lucky to avoid z^* he believes that he will be fortunate again, that is $q'(a_2,3) = 1$, then he will change his mind and choose a_2 .

Problem 5. Assume there is a finite number of income levels. An income distribution specifies the proportion of individuals at each level. Thus, an income distribution has the same methematical structure as a lottery. Consider the binary relation "one distribution is more egalitarian than another".

a. Why is the von Neumann-Morgenstern independence axiom inappropriate for characterizing this type of relation?

Assigning all members of the society the income 1 is as egalaterian as assigning all of them the income 2 and under the independence axiom, $0.5[1] \oplus 0.5[2]$ should be as egalaterian as [1], but our intuition is that $0.5[1] \oplus 0.5[2]$ is less egalitarian than assigning equal income to all members of the society.

b. Suggest and formulate a property that is appropriate, in your opinion, as an axiom for this relation. Give two examples of preference relations that satisfy this property.

If *p* and *q* are identical distributions, except that the highest (lowest) income level in *p* is less (more) than in *q*, then *p* is more egaletrian than *q*.

Example 1: $p \geq q$ if $Var(p) \leq Var(q)$.

Example 2: $p \ge q$ if $\max_{z \in \text{supp } p(z)} z - \min_{z \in \text{supp } p(z)} z \le \max_{z \in \text{supp } q(z)} z - \min_{z \in \text{supp } q(z)} z$.

Problem 6. A decision maker faces a trade-off between longevity and quality of life. His preference relation ranks lotteries on the set of all certain outcomes of the form (q,t) defined as "a life of quality q and length t " (where *q* **and** *t* **are nonnegative numbers). Assume that the preference relation satisfies von Neumann-Morgenstern assumptions and that it also satisfies the following:**

(i) There is indifference between any two certain lotteries $(q,0)$ and $[(q', 0)].$

(ii) Risk neutrality with respect to life duration: an uncertain lifetime of expected duration *T* **is equally preferred to a certain lifetime duration** *T* **when** *q* **is held fixed.**

(iii) Whatever quality of life, the longer the life the better.

a. Show that the preference relation derived from maximizing the expectation of the function $v(q)t$, where $v(q) > 0$ for all q, satisfies the **assumptions.**

(i) If $t = 0$, then $v(q)t = 0$ for all q.

(ii) Let *p* be a lottery over *t* with expectation *T*. Then

 $\sum_{i} p(t)v(q)t = v(q)\sum_{i} tp(t) = v(q)T.$

(iii) $v(q)t' > v(q)t$ for all $t' > t$.

b. Show that all preference relations satisfying the above assumptions can be represented by an expected utility function of the form $v(q)t$, where v **is a positive function.**

Since \succeq satisfies the v-NM axioms, then \succeq is represented by an expected utility function with values $w(q,t)$.

By the second property, $w(q, t)$ is a affine tranformation of *t*, that is $w(q, t) = v(q)t$ $+b(q)$.

By propery (i) it must be that $b(q) = b$ as otherwise for some q and q' we would have $w(q, 0) \neq w(q', 0)$.

By (iii) $v(q) > 0$ for all *q*.

Problem 7. Consider a decision maker who systematically calculates that $2 + 3 = 6$. Construct a "money pump" argument against him. Discuss the **argument.**

Tell the DM: "If you pay me \$5.99, I will give you two checks, one for \$2 and another for \$3." The DM will take the offer since he thinks he profits \$0.01. Then buy from him the checks for \$2.01 and \$3.01 and so on...

Problem Set 9 – Risk Aversion

Problem 1.

a. Show that a sequence of numbers (a_1, \ldots, a_K) satisfies that $\sum a_k x_k \geq 0$ for **all vectors** (x_1, \ldots, x_K) such that $x_k > 0$ for all k iff $a_k \geq 0$ for all k **.**

 \Rightarrow If *a_{k[∗]}* < 0 then consider take $x_{k^*} = 1$ and $x_k = ε > 0$ then for *ε* small enough $\sum a_k x_k < 0$.

 \leftarrow If *a_k* ≥ 0 for all *k* then $\sum a_k x_k$ ≥ 0 for all vectors $(x_1,...,x_K)$ such that x_k ≥ 0.

b. Show that a sequence of numbers (a_1, \ldots, a_K) satisfies that $\sum a_k x_k \geq 0$ for **all vectors** $(x_1,...,x_K)$ such that $x_1 > x_2 > ... x_K > x_{K+1} = 0$ iff $\sum_{k=1}^{l} a_k \geq 0$ for all *l*. It follows, like in part (a), from the equality:

 $\sum a_k x_k = \sum_{k=1}^K a_k \sum_{l=k}^K (x_l - x_{l+1}) = \sum_{l=1}^K (x_l - x_{l+1}) \sum_{k=1}^l a_k$.

Problem 2. We say that *p* **second-order stochastically dominates** *q* **and** denote this by pD_2q if $p \geq q$ for all preferences \geq satisfying the vNM **assumptions, monotonicity and risk aversion.**

a. Explain why pD_1q implies pD_2q .

If pD_1q , then $p \geq q$ for all preferences satisfying the vNM assumptions and monotonicity. Thus $p \geq q$ for all preferences satisfying the vNM assumptions, monotonicity *and* risk aversion.

b. Let p and ϵ be lotteries. Define $p + \epsilon$ to be the lottery that yields the **prize** *t* with the probability $\sum_{\alpha+\beta=\ell} p(\alpha) \epsilon(\beta)$. Interpret $p+\epsilon$. Show that if ϵ is a **lottery with expectation 0, then for all** p , $pD_2(p+\epsilon)$.

 $p + \epsilon$ is the combination of two independent lotteries p and ϵ . Let the agent satisfy vNM assumptions, monotonicity and risk aversion. Then

$$
U(p+\epsilon) = \sum_{\alpha \in \mathbb{Z}} p(\alpha) \sum_{\beta \in \mathbb{Z}} \epsilon(\beta) u(\alpha + \beta)
$$

\n
$$
\leq \sum_{\alpha \in \mathbb{Z}} p(\alpha) u(\sum_{\beta \in \mathbb{Z}} \epsilon(\beta)(\alpha + \beta)) \text{ by } u \text{ concave}
$$

\n
$$
= \sum_{\alpha \in \mathbb{Z}} p(\alpha) u(\alpha) \qquad \text{ by } Ex(\epsilon) = 0
$$

\n
$$
= U(p).
$$

c. Show that pD_2q iff for all $t < K$, $\sum_{k=0}^{t} [G(p, x_{k+1}) - G(q, x_{k+1})][x_{k+1} - x_k] \geq 0$, where $x_0 \leq \ldots \leq x_K$ are all the prizes in the support of either p or q and

 $G(p, x) = \sum_{z \geq x} p(z)$.

Let \succsim satisfy the vNM axioms, monotonicity and risk aversion. Then \succsim is represented by $U(p) = \sum_{k=0}^{K} u(x_k) p(x_k)$, with u increasing and concave. Define

$$
\alpha_k = \begin{cases} \frac{u(x_{k+1}) - u(x_k)}{x_{k+1} - x_k} & \text{if } k < K \\ 0 & \text{if } k = K \end{cases}
$$

By *u* increasing and concave, $\alpha_k \geq \alpha_{k+1}$ for all *k*. Then

$$
U(p) - U(q) = \sum_{k=0}^{K} (p(x_k) - q(x_k))u(x_k)
$$

\n
$$
= \sum_{k=0}^{K-1} [G(p, x_{k+1}) - G(q, x_{k+1})][u(x_{k+1}) - u(x_k)]
$$
 by algebra
\n
$$
= \sum_{k=0}^{K-1} [G(p, x_{k+1}) - G(q, x_{k+1})][x_{k+1} - x_k]a_k
$$
 by def. of a_k
\n
$$
= \sum_{k=0}^{K-1} [G(p, x_{k+1}) - G(q, x_{k+1})][x_{k+1} - x_k] \sum_{l=k}^{K-1} (\alpha_l - \alpha_{l+1})
$$
 telescopic sum and $\alpha_k = 0$
\n
$$
= \sum_{l=0}^{K-1} (\alpha_l - \alpha_{l+1}) \sum_{k=0}^{l} [G(p, x_{k+1}) - G(q, x_{k+1})][x_{k+1} - x_k]
$$
 by algebra
\n
$$
\geq 0
$$
 by $\alpha_l \geq \alpha_{l+1}$ and $\sum_{k=0}^{l} [G(p, x_{k+1}) - G(q, x_{k+1})][x_{k+1} - x_k] \geq 0$.

By contradiction, assume $\sum_{k=0}^{T} [G(p, x_{k+1}) - G(q, x_{k+1})][x_{k+1} - x_k] < 0$ for some *T* \lt *K*. Let $\alpha \in (0,1)$ and define

$$
u(x) = \begin{cases} x & \text{if } x \leq x_{T+1} \\ x_{T+1} + \alpha(x - x_{T+1}) & \text{if } x > x_{T+1}. \end{cases}
$$

By the above $U(p) < U(q)$ for α small enough.

Problem 3. Consider a phenomenon called preference reversal. Let $L_1 = 8/9[4] \oplus 1/9[0]$ and $L_2 = 1/9[40] \oplus 8/9[0]$.

Discuss the phenomenon that many people prefer *L*¹ **to** *L*² **but when asked to evaluate the certainty equivalence of these lotteries they attach a lower value to** L_1 **than to** L_2 .

People often prefer *L*1, but they are not willing to pay \$4 to play. Nevertheless, some people are willing to pay \$4 to play *L*2. It seems that people tend to over estimate small probabilities when they evaluate a lottery. See http://www.encyclopedia.com/doc/1O87-preferencereversal.html.

Problem 4. Consider a consumer's preference relation over *K***-tuples describing quantities of** *K* **uncertain assets. Denote the random return on the** *k***th asset by** Z_k . Assume that the random variables (Z_1, \ldots, Z_k) are **independent and take positive values with probability 1. If the consumer buys the combination of assets** (x_1, \ldots, x_K) and if the vector of realized returns **is** $(z_1,...,z_K)$, then the consumer's total wealth is $\sum_{k} x_k z_k$. Assume that the **consumer satisfies vNM assumptions, that is, there is a function** *v* **(over the sum of his returns) so that he maximizes the expected value of** *v***. Assume that** *v* **is increasing and concave. The consumer preferences over the space of the lotteries induce preferences on the space of investments. Show that the induced preferences are monotonic and convex.**

Monotonic: Let $x \geq x'$. Whenever the random variable $\sum_k x_k Z_k$ gets a certain value the random variable $\sum_k x'_k Z_k$ an higher value and thus $Ev(\sum_{k} x_{k} Z_{k}) \geq Ev(\sum_{k} x_{k}^{\prime} Z_{k}).$

Convex: Let x, x' be two investment combinations, $\lambda \in [0,1]$ and $x'' = \lambda x + (1 - \lambda)x'$. By the concavity of *v*, $v(x'' \cdot z) \geq \lambda v(x \cdot z) + (1 - \lambda)v(x' \cdot z)$ for all *z*, and thus $Ev(\sum_k x_k'' Z_k) \geq \lambda Ev(\sum_k x_k Z_k) + (1-\lambda)Ev(\sum_k x_k' Z_k)).$ The expectation of v is thus quasi-concave, and preferences are convex.

Problem 5. Adam lives in the Garden of Eden and eats only apples. Time in the garden is discrete $(t = 1, 2, \ldots)$ and apples are eaten only in discrete **units. Adam possesses preferences over the set of streams of apple consumption. Assume that:**

a) Adam likes to eat up to 2 apples a day and cannot bear to eat 3 apples a day.

b) Adam is impatient. He would be delighted to increase his consumption on day *t* **from 0 to 1 or from 1 to 2 apples at the expense of an apple he is promised a day later.**

c) In any day in which he does not have an apple, he prefers to get one apple immediately in exchange for two apples tomorrow.

d) Adam expects to live for 120 years.

Show that if (poor) Adam is offered a stream of 2 apples starting in day 4 for the rest of his expected life, he would be willing to exchange that offer for one apple right away.

```
The following is a sequence of streams, in an increasing ordering:
(0.0.0.2.2,......2)(0,0,1,0,2,\ldots,2) and continuing in this way until:
(0,0,1,1,1,......1,0)(0,0,2,0,2,0...,2,0,0)
(0,1,0,1,0,....1,0,1,0,0,0) and "folding from the end":
(0,1,0,1,0,.1,0..2,0,0,0,0,0)
(0,1,0,1,0,.1,1,0,0,0,0,0,0)...until we reach:
(0, 2, 0, \ldots, 0)(1,0,...)
```
Problem 6. In this problem you will encounter Quiggin and Yaari's functional, one of the main alternatives to expected utility theory.

 \boldsymbol{Recall} that expected utility \boldsymbol{can} be written as $U(p) = \sum_{k=1}^K p(z_k) u(z_k)$ where $z_0 < z_1 < ... < z_K$ are the prizes in the support of p . Let *W*(*p*) = $\sum_{k=1}^{K} f(G_p(z_k))[z_k - z_{k-1}]$, where *f* : [0,1] → [0,1] is a continuous increasing function and $G_p(z_k) = \sum_{i \geq k} p(z_j)$. ($p(z)$ is the probability that the lottery p yields z and G_p is the "anti-distribution" of p **.**)

a. The literature often refers to *W* **as the dual expected utility operator. In what sense is** *W* **dual to** *U***?**

 $\text{Recall that } Ex(p) = \sum_{k=1}^{K} p(z_k) z_k = \sum_{k=1}^{K} G_p(z_k) [z_k - z_{k-1}]$

While the expected utility functional transforms the prize numbers whereas Quiggin-Yaari functional transofrms the anti-distribution numbers.

b. Show that *W* **induces a preference relation on** *LZ* **which satisfies the continuity axiom but may not satisfy the independence axiom.**

Preferences are continuous because *U* is clearly continuous.

Let $K = 2, f(x) = x^2, z_0 = 0, z_1 = 1$ and $z_2 = 4$. Define lotteries $p = 75[z_0] \oplus 25[z_2]$ and $p' = 5[z_0] \oplus 5[z_1]$. Then

$$
U(p) = 4f(\frac{1}{4}) = \frac{1}{4} = f(\frac{1}{2}) = U(p')
$$

but

 $U(\frac{1}{2}p \oplus \frac{1}{2}[z_1]) = f(\frac{5}{8}) + 3f(\frac{1}{8}) = \frac{28}{64} < \frac{9}{16} = f(\frac{3}{4}) = U(\frac{1}{2}p' \oplus \frac{1}{2}[z_1]).$

c. What are the difficulties with a functional form of the type $\sum_{z} f(p(z))u(z)$? **(See Handa (1977))**

(1) If the DM is indifferent between prizes z_1 and z_2 , then $[z_1]$ and $0.5[z_1] \oplus 0.5[z_2]$ need not be indifferent. If $f(1/2) \neq 1/2$, then $U([z_1]) \neq U(0.5[z_1] \oplus 0.5[z_2]).$

(2) The DM might be "worse" off if probability weight is shifted to a more preferred alternative:

if $f(1/2)$ > 1/2, 0.5[1 – ε] \oplus 0.5[1] ≻ [1] for ε > 0 small enough while and– if $f(1/2) < 1/2$, $0.5[1 + \varepsilon] \oplus 0.5[1] < [1]$ for $\varepsilon > 0$ small enough.
Problem 7. The two envelopes paradox

Assume that a number 2*ⁿ* **is chosen with probability** 2*ⁿ*/3*ⁿ*¹ **and the amounts of money** 2*ⁿ***,** 2*ⁿ*¹ **are put into two envelopes. One envelope is chosen randomly and given to you and the other is given to your friend. Whatever the amount of money in your envelope, the expected amount in your friend's envelope is larger (verify it). Thus, it is worthwhile for you to switch envelopes with him even without opening the envelope! What do you think about this paradoxical conclusion?**

Note that this is indeed a probability distribution:

 $\sum_{n=0}^{\infty} \frac{2^n}{3^{n+1}} = \frac{1}{3} \sum_{n=0}^{\infty} \left(\frac{2}{3}\right)$ $\frac{2}{3}$)ⁿ = $\frac{1}{3}$ 1 $\frac{1}{1-\frac{2}{3}}=1.$

Assume that in your envelope the sum of money is 2^n . For any $n \geq 1$ this can be either the smaller amount or the larger one. If it is the smaller, then the other envelope contains 2^{n+1} and changing envelopes means a gain of 2^n . The probability for this event is $2^n/3^{n+1}$. If your amount is the larger one, then the other envelope contains 2*ⁿ*−¹ and changing means a loss of 2*ⁿ*−¹. The probability for this event is 2^{*n*−1}/3^{*n*}. Hence, the expected gain when changing envelopes is $\frac{2^n \cdot 2^n/3^{n+1}-2^{n-1}\cdot 2^{n-1}/3^n}{2^n/3^{n+1}+2^{n-1}/3^n} = \frac{2^n}{10}$ which is positive for any *n*. For *n* = 0, your envelope is surely the smaller one, hence changing envelopes is profitable for any $n \geq 0$. Thus, we

can conclude that you should change envelopes without even opening yours.

Note that the random expected amount of money in any envelope, your own and the other one, is $\frac{1}{3} + \sum_{n=1}^{\infty} 2^n \cdot 5 \cdot \frac{2^{n-1}}{3^{n+1}} = \infty$. It is possible to show that if the problem is constructed such that the expected value of each envelope is finite, this paradox does not arise.

Random variables with infinite expectation create many paradoxes. For example, after every draw of such a random variable the decision maker who is risk neutral would prefer to replace the outcome in hand with another draw...

Problem Set 10 – Social Choice

1. Assume that the set of social alternatives, *X***, includes only two alternatives. Define a social welfare function to be a function that attaches a preference to any profile of preferences (allow indifference for the SWF and the individuals' preference relations). Consider the following axioms:**

Anonymity: If σ is a permutation of N and if $p = \{\succsim_i\}_{i\in N}$ and $p' = \{\succsim'_i\}_{i\in N}$ are \mathbf{t} wo profiles of preferences on X so that $\succsim'_{\sigma(i)} \;\; = \succsim_i$, then $\succsim (p) \; = \succsim (p').$

Neutrality: For any preference \succsim _{*i*} , define $(-\succeq_i)$ as the preference **satisfying** $x(-\succcurlyeq_i y)$ iff $y \succcurlyeq_i x$. Then, $\succcurlyeq_i \left\{ \frac{1}{2} \right\} \succcurlyeq_i \left\{ \frac{1}{2} \right\} \succcurlyeq_i \left\{ \frac{1}{2} \right\}$.

Positive Responsiveness: If the profile *ⁱ* ′ *ⁱ*∈*^N* **is identical to** *ⁱ ⁱ*∈*^N* **with** the exception that for one individual j either $(x\sim_j y$ and $x\succ'_j y$) or $(y\succ_j x$ and $(x \sim'_j y)$ and if $x \succcurlyeq y$, then $x \succ' y$.

a. Interpret the axioms.

A: The social aggregation treats any two individuals symmetrically.

N: The social aggregation treats the two alternatives symmetrically.

PR: The fact that one individual changed his mind in favor of an alternative cannot harm (and in some cases is required to improve) the social status of that alternative.

b. Show that majority rule satisfies all of them.

A: trivially satisfied.

N: Let $N(x, p)$ be the number of individuals that strictly prefer x to y in profile p. If $x \geq (p)y$, then $N(x, p) \geq N(y, p)$, and thus $y \geq (-p)x$, proving N.

PR: *p*′ is identical to *p*, except that one individual "increases" his preference for *x*, and $N(x,p) \geq N(y,p)$ then $N(x,p') > N(y,p')$, and thus $x \succ (p')y$.

c. Prove May's theorem by which the majority rule is the only SWF satisfying the above axioms.

Assume $N(x,p) = N(y,p)$. Let σ be a permutation such that $\sigma(i)$'s preference is the reverse of *i*'s preference. Let $p'=\{\succsim'_i\}_{i\in N}$ be a profile so that $\succsim'_{\sigma(i)}=\succsim_i$. By A $(x \geq (p)y$ iff $x \geq (p')y$. Since $p' = -p$ by N $x \geq (p')y$ iff $y \geq (p)x$ and thus $x \sim (p)y$.

Assume *p* a profile such that $N(x, p) > N(y, p)$. Assume $y \ge (p)x$. We can move from *p* to a profile *p'* by changing the preferences of $N(x,p) - N(y,p)$ individuals who prefer x to y to indifference. By PR we would get $y \succ (p')x$ although $N(x, p') = N(y, p')$, a contradiction.

d. Are the above three axioms independent?

Yes.

A and N, but not PR: $x \sim (p)y$ for all p.

A and PR, but not N: Let $X = \{a,b\}$. $a > (p)b$ if $N(a,p) > N(b,p) + 1$, otherwise $b \succ (p)a$.

N and PR, but not A: For any profile of preferences we attach the

"lexicographic" preferences with some fixed priority of the individuals (For example, $x \geq (p)y$ if $[x \succ_1 y]$ or $[x \sim_1 y]$ and $x \succ_2 y$] and so on...)

2. Assume that the set of alternatives, X , is the interval $[0,1]$ and that each **individual's preference is single-peaked, i.e., for each** *i* **there is an alternative** a_i^* such that if $a_i^* \geq b > c$ or $c > b \geq a_i^*$, then $b \succ_i c$. Show that for any odd n , if **we restrict the domain of preferences to single-peaked preferences, then the majority rule induces a "well-behaved" SWF.**

Let $x \geq p$ *y* and $y \geq p$ *z*. By contradiction, assume that $z \geq p$ *x*.

(i) $x < y < z$. If a majority find x as good as y then all those strictly prefer y to z and there is a majority who strictly prefer *x* over *z*.

(i) $x < z < y$. If a majority find *y* as good as *z* then all those strictly prefer *z* to *x* and there is a majority who strictly prefer *y* over *x*.

(i) $z < x < y$. If a majority find z as good as x then all those strictly prefer x to y and there is a majority who strictly prefer *z* over *y*.

3. Each of *N* **individuals chooses a single object from among a set** *X***, interpreted as his recommendation for the social action. We are interested in functions that aggregate the individuals' recommendations (not preferences, just recommendations!) into a social decision (i.e.,** $F: X^N \rightarrow X$). Discuss the **following axioms:**

Par: If all individuals recommend *x*[∗], **then the society chooses** *x*[∗]**.**

I: If the same individuals support an alternative *x* ∈ *X* **in two profiles of recommendations, then** *x* **is chosen in one profile if and only if it chosen in the other.**

a. Show that if *X* **includes at least three elements, then the only aggregation method that satisfies** *P* **and** *I* **is a dictatorship.**

L1: For any recommendation profile $(x_1,...,x_N)$, $F(x_1,...,x_N) \in \{x_1,...,x_N\}$. Otherwise, by *I*, $F(c_1, c_2) = F(x_1, \ldots, x_N)$ for any $c \neq F(x_1, \ldots, x_N)$, contradicting *P*. **L2**: If $F(x_1,...,x_N) = a$ and for all $G = \{i | x_i = a\} \subset \{i | y_i = a\}$ then $F(v_1,\ldots,v_N)=a.$

If not than by L1 $F(y_1,..., y_N) = b \neq a$ where *b* is one of the alternatives in *y*.

Let *c* be a third alternative. Let $(z_1,...,z_N)$ be the same as *y* with any $i \notin G$ with $y_i = a$ is changed to $z_i = c$. Then, $\{i | y_i = b\} = \{i | z_i = b\}$ and $\{i | x_i = a\} = \{i | z_i = a\}$ thus by \mid *F* (z) should be both *b* and *a*. A contradiction.

Let us say that *G* is *decisive* with respect to $x^* \in X$, if [for all $i \in G$, $x_i = x^*$] then $[F(x_1,...,x_N) = x^*].$

By L2 if $F(x_1,...,x_N) = a$ then $\{i | x_i = a\}$ is decisive with respect to a.

L3: If *G* is decisive with respect to *a* with $|G|$ \geq 2, then for any *b* there exists $\emptyset \subset G' \subset G$ such that G' is decisive with respect to a or b.

Let *c* be a third alternative. Since *G* is decisive with respect to *a* then $F(x_1,...,x_N) = a$ where $x_i = a$ for $i \in G$ and $x_i = c$ otherwise

Let G_1 , G_2 be a partition of G and

$$
y_i = \begin{cases} a & \text{if } i \in G_1 \\ b & \text{if } i \in G_2 \\ c & \text{if } i \in N \setminus G \end{cases}
$$

By $IF(y_1,..., y_N)$ is not *c* and by L1 it is either *a* or *b*. Thus, by L2 either G_1 or G_2 are decisive with respect to *a* or *b*.

L4: There is a singleton *i* [∗] who is decisive with respect to some alterative *a*.

L5: *i* [∗] is decisive regarding any alterative *b*.

Let $x_i^* = a$ and $x_i = c$ for all other *i*. By I, $F(x) = a$.

Let $y_{i^*} = b$ and $y_i = c$ for all other *i*, By $\mid F(y) \neq c$ and by L1 $F(y) = b$ and by L2 it is decisive regarding *b*.

b. Show the necessity of the three conditions P , I , and $|X|\geq 3$ for this **conclusion.**

Choosing the most popular recommendation (with pre-specified tie breaking rule) satisfies P and I when $|X|=2$.

When $|X|\geq 3$, the most popular recommendation (with pre-specified tie breaking

rule)satisfies *P* but fails *I*.

Always choosing action $x^* \in X$, regardless of recommendation, satisfies *I* but fails *P*.

4. Who is an economist? Departments of economics are often sharply divided over this question. Investigate the approach according to which the determination of who is an economist is treated as an aggregation of the views held by department members on this question.

Let $N = \{1, ..., n\}$ be a group of individuals ($n \geq 3$). Each $i \in N$ "submits" a **set** *Ei***, a proper non empty subset of** *N***, which is interpreted as the set of "real economists" in his view. An aggregation method** *F* **is a function that assigns a proper non empty subset of** *N* to each profile $(E_i)_{i=1,\dots,n}$ of proper subsets of *N***.** $F(E_1, \ldots, E_n)$ is interpreted as the set of all members of *N* who **are considered by the group to be economists. (Note that we require that all opinions be proper subsets of** *N***.) Consider the following axioms on** *F***:**

Consensus: If $j \in E_i$ for all $i \in N$, then $j \in F(E_1, \ldots, E_n)$ and if $j \notin E_i$ for all $i \in N$, then $j \notin F(E_1, \ldots, E_n)$.

Independence: If (E_1, \ldots, E_n) and (G_1, \ldots, G_n) are two profiles of views so **that for all** $i \in N$, $j \in E_i$ iff $j \in G_i$, then $j \in F(E_1, \ldots, E_n)$ iff $j \in F(G_1, \ldots, G_n)$.

a. Interpret the two axioms.

C: If all individuals include (omit) an economist from their list, then that option is included (omitted) from the aggregation.

I: If all individuals in profiles $\{E_i\}$ and $\{G_i\}$ have the same opinion regarding *j*, then *j* is either included or excluded in both aggregations.

b. Find one aggregation method that satisfies *C* **but not** *I* **and one that satisfies** *I* **but not** *C***.**

Select the set with the smallest number of elements suggested by one of the members (for tie breaking rule, from among the sets with minimal number of individuals choose the one suggested by the member with the smallest index) This methods satisfies *C* but not *I*.

Always selecting $F(E_1,...,E_n) = \{1\}$ satisfies *I* but not *C*.

c. Provide a proof similar to that of Arrow's Impossibility Theorem of the claim that the only aggregation methods that satisfy the above two axioms are those for which there is a member *i*^{*} such that $F(E_1,...,E_n) = E_{i^*}$.

L1: Assume that *G* is almost decisive regarding *j* (that is *j* is an economist whenever the group of people who consider *j* an economist is precisely *G*) then it is almost decisive regarding any *h* (and thus will be almost decisive).

Consider the profile where the supporters of *j* are the members of *G*, the supporters of *i* (a third member) are *N* − *G* and everybody supports all other members. By C those must be in *E* and since *E* is a proper subset and $j \in E$, it must be that *i* is not.

Now consider the profile where the set of supporters of *h* is *G*, the supporters of *i* are *N* − *G* and nobody considers any other member to be an economist. By C all those members besides *i* and *h* are not in *E*. By I *i* is not in *E*. Since *E* is a proper subset it must be that *h* is in *E*. Thus, *G* is almost decisive regarding *h*.

L2: If G is almost decisive and contains more than one element then there is a

proper subset of *G* which is almost decisive.

Partition *G* to G_1 and G_2 . Consider the profile where 1 is considered an economist by exactly the members of *G*1, 2 is considered an economist by exactly the members of G_2 and all other individuals are considered economists by the members of *N* − *G* only. As we have seen in L1, since *G* is almost decisive then *N* − *G* is not and thus all individuals besides 1 and 2 are not determined by the aggregator to be economists. It follows that either 1 or 2 must be an economist and thus (using I) either G_1 or G_2 is almost decisive in regard 1 and 2 and by L1 at least one of them is almost decisive.

L3: There is an *i*^{*} who is almost decisive.

L4: If *i* is supported by G which contains i^* then i is an economist.

Consider the profile where *i* is supported by *G*, *j* by $N - \{i^*\}$ and all the rest by no one. By C all members $N - \{i^*, j\}$ are not economists. From the argument above neither is *j* so *i* must be an economist with respect to this profile and by I with respect to any profile.

Review Problems

A. Choice:

Problem A1 (Princeton 1997)

A decision maker forms preferences over the set X of all possible distributions of a population over two categories (such as living in two locations). An element in X is a vector (x_1, x_2) where $x_i \geq 0$ and $x_1 + x_2 = 1$. The decision maker has two *considerations in mind:*

He thinks that if $x \geq y$ *, then for any z, the mixture of* $\alpha \in [0,1]$ *of x with* $(1 - \alpha)$ *of z should be at least as good as the mixture of* α of γ *with* $(1 - \alpha)$ of z *.*

He is indifferent between a distribution that is fully concentrated in location 1 and one that is fully concentrated in location 2.

a. Show that the only preference relation that is consistent with the two principles is the degenerate indifference relation $(x \sim y)$ *for any* $x, y \in X$ *).*

Let $x = (1,0), y = (0,1)$ and $(\alpha,1 - \alpha)$ be an arbitrary distribution, $0 \le \alpha \le 1$. By the second consideration, $x \sim y$, and in particular $x \ge y$. By the first consideration, the mixture α of x and $(1 - \alpha)$ of y is at least as good as the mixture α of y and $(1 - \alpha)$ of *y* (choosing *z* to be *y*). Thus, $(\alpha, 1 - \alpha) \ge y$. Similarly, the mixture α of *x* and $(1 - \alpha)$ of *x* is at least as good as the mixture α of *x* and $(1 - \alpha)$ of *y* (choosing *z* to be *x*). Thus, $x \geq (\alpha, 1 - \alpha)$. Hence, any distribution $(\alpha, 1 - \alpha)$ is indiffernet to *x* and *y*, since $x \geq (\alpha, 1 - \alpha) \geq y$.

b. The decision maker claims that you are wrong because his preference relation is represented by a utility function $|x_1 - 1/2|$ *. Why is he wrong?*

The preferences represented by the utility function $u(x_1, x_2) = |x_1 - \frac{1}{2}|$ satisfy the first consideration, since $u(1,0) = |1 - \frac{1}{2}| = |0 - \frac{1}{2}| = u(0,1)$. However, these preferences fail to satisfy the second consideration: although $(1,0) \geq (0,1)$, mixing both sides with $\alpha = \frac{1}{2}$ of $(0, 1)$ yields $(\frac{1}{2}, \frac{1}{2}) \prec (0, 1)$, since $u(\frac{1}{2}, \frac{1}{2}) =$ $\left|\frac{1}{2} - \frac{1}{2}\right| < |0 - \frac{1}{2}| = u(0,1).$

Problem A2 (Princeton 2000. Based on Fishburn and Rubinstein (1982).)

Let $X = \mathbb{R}^+ \times \{0, 1, 2, \dots\}$, where (x, t) is interpreted as receiving $\& x$ at time *t*. A *preference relation on X has the following properties:*

(i) *There is indifference between receiving* \$0 *at time* 0 *and receiving* \$0 *at any other time.*

(ii) *It is better to receive any positive amount of money as soon as possible.*

(iii) *Money is desirable.*

(iv) The preference between (x, t) and $(y, t + 1)$ is independent of t.

(v) *Continuity.*

a. Define formally the continuity assumption for this context.

For any (x, t) and (y, s) with $(x, t) \succ (y, s)$, there is $\varepsilon > 0$ such that for any x' such that $|x - x'| < \varepsilon$ and any y' such that $|y - y'| < \varepsilon$, $(x', t) > (y', s)$.

b. Show that the preference relation has a utility representation.

Claim 1: For any pair (x, t) , there is a unique number $v(x, t) \in \Re$ such that $(x,t) \sim (v(x,t),0).$

Proof: By (iii) it is enough to show that for every (x, t) there is y with $(x, t) \sim (y, 0).$

For $x = 0$, by (i) $(0, t) \sim (0, 0)$.

For any pair (x, t) with $x \neq 0$, define the set $B(x, t) = \{z \in \Re_+(x, t) \geq (z, 0)\}$. By (i) and (iii), $0 \in B(x,t)$. By (ii) and (iii) $B(x,t)$ is bounded above by x. Let

 $x^* = \sup B(x,t)$. Then by continuity $(x,t) \sim (x^*,0)$.

Claim 2: The preference relation is represented by $u(x,t) = v(x,t)$. **Proof:**

$$
u(x,t) \ge u(y,s) \iff v(x,t) \ge v(y,s)
$$

\n
$$
\iff (v(x,t),0) \ge (v(y,s),0) \text{ (by (ii))}
$$

\n
$$
\iff (x,t) \ge (y,s) \text{ (by def of } v \text{. and transitivity)}
$$

c. Verify that the preference relation represented by the utility function $v(x,t) = u(x)\delta^t$ (with $\delta < 1$ and *u* continuous, increasing and $u(0) = 0$) satisfies the *above properties.*

(i): For all *t*, *t'*, $v(0,t) = u(0)\delta^t = 0 = u(0)\delta^s = v(0,s)$. Thus, $(0,t) \sim (0,s)$.

(ii): For any *x* and $t < s$, $v(x,t) = \delta^t u(x) > \delta^s u(x) = v(x,s)$. Thus, $(x,t) > (x,s)$.

(iii): This holds since $u(\cdot)$ is increasing.

(iv): For all x, y, t, s ,

$$
(x,t) \ge (y, t+1) \iff \delta^t u(x) > \delta^{t+1} u(y)
$$

$$
\iff \delta^s u(x) > \delta^{s+1} u(y)
$$

$$
\iff (x,s) \ge (y,s+1)
$$

(v): Continuity follows from *u* being continuous.

d. Formulize a concept "one preference relation is more impatient than another".

 \geq ¹ is more impatient than \geq ² if for any (x, t) , and any (y, s) with $s > t$, $y > x$ $(y,s) \geq^1 (x,t) \Rightarrow (y,s) \geq^2 (x,t).$

e. Discuss the claim that preferences represented by $u_1(x)\delta_1^t$ are more impatient *than preferences represented by* $u_2(x)\delta_2^t$ *if and only if* $\delta_1 < \delta_2$.

This would be true when $u_1 = u_2 = u$: then for any $s > t$, $y > x$, $\delta_1^s u(y) \geq \delta_1^t u(x) \Leftrightarrow \delta_1^{(s-t)} \geq u(x)/u(y)$ implying $\delta_2^{(s-t)} \geq u(x)/u(y) \Leftrightarrow \delta_2^s u(y) \geq \delta_2^t u(x)$

However, the claim is not necessarily true if $u_1 \neq u_2$. For instance, suppose $u_1(x) = \ln x$, $u_2(x) = x$, $\delta_1 = 1/2$ and $\delta_2 = 1/\sqrt{3} > 1/2$. Then person 2 is indifferent between getting \$1 at time 0 and \$3 at time 2, since

$$
\delta_2^2 u_2(3) = 1 = \delta_2^0 u_2(1)
$$

However, person 1, the supposedly more impatient (lower δ) person, prefers to wait

for the \$3:

$$
\delta_1^2 u_1(3) = 1/4 \ln 3 > \delta_1^0 u_1(1) = \ln 1.
$$

Problem A3 (NYU 2005, inspired by Chen, M.K., V. Lakshminarayanan and L. Santos (2005).)

In an experiment, a monkey is given $m = 12$ coins which he can exchange for *apples or bananas. The monkey faces m consecutive choices in which he gives a coin either to an experimenter holding a apples or another experimenter holding b bananas.*

1. *Assume that the experiment is repeated with different values of a and b and that each time the monkey trades the first* 4 *coins for apples and the next* 8 *coins for bananas.*

Show that the monkey's behavior is consistent with the classical assumptions of consumer behavior (namely, that his behavior can be explained as the maximization of a monotonic, continuous and convex preference relation on the space of bundles).

We can model the problem of the monkey as the following. It has wealth *m* and can buy two commodities, apples and bananas. The price of one apple is $p_a = 1/a$ and the price of one banana is $p_b = 1/b$. What the experiment shows is that the monkey always spends 1/3 of its wealth on apples and 2/3 on bananas. This behavior is consistent with Cobb-Douglas preferences with weights 1/3 and 2/3.

2. *Assume that it was later observed that when the monkey holds an arbitrary number of coins, m. Then, irrespective of the values of a and b, he exchanges the first* 4 *coins for apples and the remaining m* − 4 *coins for bananas. Is this behavior consistent with the rational consumer model?*

No. Assume, by contradiction, that \geq is monotonic and rationalizes the monkey behavior.

Let $m = 4$, $a = 2$ and $b = 4$. In this case the monkey would buy the following bundle $(8,0)$. Since the monkey could also buy $(4,8)$, it must be the case that $(8,0) \ge (4,8)$. Monotonicity of \ge implies that $(9,1) \ge (4,8) \ge (4,6)$.

Now let's look at the monkey's choice when $m = 10$, $a = 1$ and $b = 1$. In this case the monkey would chose $(4,6)$ though $(9,1)$ is feasible. A contradiction.

Problem A4 (NYU 2005)

Let X be a finite set containing at least three elements. Let C be a choice correspondence. Consider the following axiom:

If $A, B \subseteq X, B \subseteq A$ and $C(A) \cap B \neq \emptyset$, then $C(B) = C(A) \cap B$.

1. *Show that the axiom is equivalent to the existence of a preference relation such that* $C(A) = \{x \in A \mid x \succ a$ *for all* $a \in A\}$.

It's enough to show that the Choice Axiom in the question is equivalent to the

Weak Axiom.

Choice Axoim \Rightarrow WA. Let $x \in C(A)$, $y \in C(B)$ and $x, y \in A \cap B$. Define $B' = A \cap B$. Clearly, $C(A) \cap B' \neq \emptyset$ and $C(B) \cap B' \neq \emptyset$, so the Choice Axiom implies that *C*(*A*) ∩ *B* = *C*(*A*) ∩ *B*['] = *C*(*B*[']) = *C*(*B*) ∩ *B*['] = *C*(*B*) ∩ *A*. But *x* clearly is in *C*(*A*) ∩ *B*, which implies that $x \in C(B) \cap A$, which itself implies that $x \in C(B)$.

WA ⇒ Choice Axiom. Let *B* ⊆ *A* and $C(A) \cap B \neq \emptyset$. Let $x \in C(B)$ and *y* ∈ *C*(*A*) ∩ *B*. Clearly, *x*, *y* ∈ *A* ∩ *B*. But then the WA implies that *x* ∈ *C*(*A*), and therefore, $x \in C(A) \cap B$ and $y \in C(B)$. But this means that $C(B) \subseteq C(A) \cap B$ and $C(A) \cap B \subseteq C(B)$, which is equivalent to $C(B) = C(A) \cap B$.

2. *Consider a weaker axiom:*

If $A, B \subseteq X, B \subseteq A$ and $C(A) \cap B \neq \emptyset$, then $C(B) \subseteq C(A) \cap B$. *Is this sufficient for the above equivalence?*

No, it's not sufficient. Consider the following example. Let $X = \{a, b, c\}$. Suppose $C(X) = X$, $C({a,b}) = {a}$, $C({b,c}) = {b}$, $C({a,c}) = {c}$. If there was a preference \succeq such that $C(A) = \{x \in A \mid x \succ a$ for all $a \in A\}$ then $a \succ b$, $b \succ c$ and $c > a$, a contradiction.

Problem A5 (NYU 2006)

Consider a world with balls of K different colors. An object is called a bag and is specified by a vector $x = (x_1, \ldots, x_K)$ (where x_k is a non-negative integer indicating *the number of balls of color k). For convenience denote by* $n(x) = \sum x_k$ *the number of balls in bag x.*

An individual has a preference relation over bags of balls.

1. *Suggest a context where it will make sense to assume that:*

i. For any integer λ , $x \sim \lambda x$.

ii. If $n(x) = n(y)$, then $x \ge y$ iff $x + z \ge y + z$.

Suppose that the decision maker chooses a ball from bag *x*, and he gets a prize depending on the color of the ball he chose. Then we can identify a bag *x* with a lottery $p(x)$ in which the prize associated with the color k is received with probability $x_k/n(x)$. Since $p(x) = p(\lambda x)$, we should expect $x \sim \lambda x$. Moreover, *ii* would follow in this case from the independence axiom.

2. *Show that any preference relation which is represented by* $U(x) = \sum x_k v_k/n(x)$ for some vector of numbers (v_1, \ldots, v_k) satisfies the two axioms. A1: For any natural number λ ,

$$
U(\lambda x) = \sum_{k} \frac{\lambda x_k}{n(\lambda x)} v_k = \sum_{k} \frac{\lambda x_k}{\lambda n(x)} v_k = U(x).
$$

A2: If $n(x) = n(y)$, then

$$
U(x) \ge U(y) \iff \sum_{k} x_{k}v_{k} \ge \sum_{k} y_{k}v_{k}
$$

$$
\iff \sum_{k} (x_{k} + z_{k})v_{k} \ge \sum_{k} (y_{k} + z_{k})v_{k}
$$

$$
\iff U(x + z) \ge U(y + z)
$$

where in the last equivalence we use that fact that $n(x+z) = n(y+z)$.

3. *Find a preference relation which satisfies the two properties that cannot be represented in the form suggested in (2).*

Let $K = 3$ and let \geq_L be the usual lexicographic relation on \mathfrak{R}^3 . Define the preference relation \succsim on *X* as

$$
x \succeq y \Leftrightarrow (x_1/n(x), x_2/n(x), x_3/n(x)) \succeq_L (y_1/n(y), y_2/n(y), y_3/n(y)).
$$

A1: Note that for any *x* and any natural number λ , $x_i/n(x) = \lambda x_i/\lambda n(x)$ for all *i*. Hence, $x \sim \lambda x$.

A2: $x \ge y$ and $n(x) = n(y)$ iff $(x_1, x_2, x_3) \ge L (y_1, y_2, y_3)$ iff $(x_1 + z_1, x_2 + z_2, x_3 + z_3) \geq L (y_1 + z_1, y_2 + z_2, y_3 + z_3)$ iff $x + z \geq y + z$ (since $n(x + z) = n(y + z)$.

To see that \geq does not admit a representation of the form given in part (2), suppose to the contrary that $U(x)$ represents \geq . Note that $(1,0,n) > (0,1,0)$ for all *n*. Hence, $v_1/(n+1) + v_3 n/(n+1) > v_2$ for all *n*. Passing to limit as $n \to \infty$ gives $v_3 \ge v_2$. On the other hand, $(0,1,0)$ \geq $(0,0,1)$ implies $v_2 > v_3$, a contradiction.

Problem A6 (NYU 2007. Based on Plott (1973).)

Let X be a set and C be a choice correspondence defined on all non-empty subsets of X.

We say that C satisfies Path Independence (PI) if for every two disjoint sets A and B, we have $C(A \cup B) = C(C(A) \cup C(B))$ *.*

We say that C satisfies Extension (E) if $x \in A$ *and* $x \in C(\{x, y\})$ for every $y \in A$ *implies that* $x \in C(A)$ *for all sets* A *.*

1. *Interpret PI and E.*

PI: The agent's choice set is identical, regardless if he (1) chooses directly from the set *A* or (2) first partitions *A* into two subsets, chooses from each of the subsets and then makes a subsequent choice from these two choice sets.

E: If $x \in A$ is chosen when compared (pairwise) with every other alternative in A, then *x* is in the choice set of *A*.

2. *Show that if C satisfies both PI and E, then there exists a binary relation that is complete, reflexive and satisfies* $x \succ y$ *and* $y \succ z$ *implies* $x \succ z$ *, such that* $C(A) = \{x \in A \mid \text{for no } y \in A \text{ is } y \succ x\}.$

For all $x, y \in X$, define $x \ge y$ if $x \in C({x, y})$. Clearly, \ge is complete and reflexive. By contradiction, assume $x \succ y$ and $y \succ z$, but $z \succ z$, that is, $C(\lbrace x, y \rbrace) = \lbrace x \rbrace$, $C({y,z}) = {y}$, and $z \in C({x,z})$. Define $A = {x,y,z}$, and note that PI implies that

 $C(A) = C(C(Jx, y), 1 + C(Jz)) = C(Jx, z) = C(C(x, z))$

$$
C(A) = C(C(\langle x, y \rangle) \cup C(\langle z \rangle)) = C(\langle x, z \rangle) \Rightarrow z \in C(A) \text{ and}
$$

 $C(A) = C(C(\lbrace x \rbrace) \cup C(\lbrace y, z \rbrace)) = C(\lbrace x, y \rbrace) \Rightarrow z \notin C(A),$

a contradiction. Therefore, $x > z$.

Finally, if $x \ge y$ for all $y \in A$, then $x \in C(A)$ by E, and thus $C(A) \supseteq \{x \in A \mid \text{ for no } y \in A, y \succ x\}$. Conversely, if there exists a $y \in A$ such that $y \succ x$, then by PI

$$
C(A) = C(C(\lbrace x,y \rbrace) \cup C(A \setminus \lbrace x,y \rbrace)) = C(\lbrace y \rbrace \cup C(A \setminus \lbrace x,y \rbrace)),
$$

and thus $x \notin C(A)$. Thus, $C(A) = \{x \in A \mid \text{for no } y \in A, y \succ x\}.$

3. *Give one example of a choice correspondence satisfying PI but not E, and one satisfying E but not PI.*

Let \succeq be an ordering over X.

PI and not E: Define $C(A)$ to equal the best and second best elements in A according to \succeq .

Satisfies PI: The best and second best elements of $A \cup B$ are contained in $C(A) \cup C(B)$, and thus $C(C(A) \cup C(B))$ equals these first and second best elements in $A \cup B$.

Fails E: Every $x \in A$ is in $C(\{x, y\})$ for all $y \in A$. Nevertheless, only the two best elements in *A* are included in *CA*.

PI and not E: Define $C(A)$ to be the best element in *A* if $A \neq X$, and $C(X) = X$. **Satisfies E:** If $x \in C(\{x, y\})$ for all $y \in A$, then there is no $y \succ x$, and thus

 $x \in C(A)$.

Fails PI: Let *A* and *B* be a partition of *X*. Then $C(X) \neq C(C(A) \cup C(B))$.

Problem A7 (NYU 2008) (based on work of Kfir Eliaz, Michael Richter, and Ariel Rubinstein)

Let X be a (finite) set of alternatives. Given any choice problem *A* (where $|A| \ge 2$), *the decision maker chooses a set* $D(A) \subseteq A$ *of two alternatives which he wants to examine more carefully before making the final decision.*

The following are two properties of D:

A1: If $a \in D(A)$ and $a \in B \subset A$ then $a \in D(B)$. *A2: If* $D(A) = \{x, y\}$ *and* $a \in D(A - \{x\})$ for some *a* different than *x* and *y*, then $a \in D(A - \{y\})$.

Answer the following four questions. A full proof is required only for the last question:

1. *Find an example of a D function which satisfies both A1 and A2.*

Let $>$ be a strict preference relation on X. Let $D(A)$ be the set of the two \succ -best elements in *A*.

2. *Find a function D which satisfies A1 and not A2.*

Let $>$ be a strict preference relation on X. Let $D(A)$ be the set containing the \succ -best element and the \succ -worst element in A.

3. *Find a function D which satisfies A2 and not A1.*

Let $>$ be a strict preference relation on X. Let $D(A)$ be the set containing the second and third \succ -best elements in *A* (if $|A|\geq 3$) and $D(A) = A$ otherwise.

4. *Show that for any function D satisfying A1 and A2 there exists an ordering of the elements of X s.t.* $D(A)$ *is the set of the two* \rightarrow -best elements in A *.*

Proof: Let *D* be a choice correspondence satisfying the three axioms. Let us construct inductively a count of all elements in *X*.

Let $D(X) = \{x_1, x_2\}$.

Assume we constructed the sequence $x_1, x_2, \ldots, x_K, x_{K+1}, x_{K+2}$ such that *D*(*X* – $\{x_1, x_2, \ldots, x_k\}$) = $\{x_{k+1}, x_{k+2}\}$ for all $0 \le k \le K$.

By *A*1 x_{K+2} ∈ $D(X - \{x_1, x_2, \ldots, x_K, x_{K+1}\})$. Define x_{K+3} by

D(*X* – $\{x_1, x_2, \ldots, x_{K+1}\}$) = $\{x_{K+2}, x_{K+3}\}$.

Define $x_i > x_j$ if $i < j$.

For any set *A* let $x_{i(A)}$ and $x_{i(A)}$ be the two \rightarrow -top elements in *A*. The set $D(A)$ must contain the \succ -top element in Λ since by definition

 $D(X - \{x_1, \ldots, x_{i(A)-1}\}) = \{x_{i(A)}, x_{i(A)+1}\}$ and as $A \subseteq X - \{x_1, \ldots, x_{i(A)-1}\}$ then by $A1$ $x_{i(A)} \in D(A)$.

Assume by contradiction that there exists a set A such that $D(A)$ contains $x_{i(A)}$, but not $x_{i(A)}$. Consider such a set *A* with minimum $i(A) - i(A)$. We will show that $j(A) - i(A) = 1$ which leads to contradiction (since by construction *D*($\{x_k | k \ge i(A)\}$) = $\{x_{i(A)}, x_{i(A)+1}\}$ and by *A*1 *D*(*A*) = $\{x_{i(A)}, x_{i(A)+1}\}$.

If *j*(*A*) > *i*(*A*) + 1 consider the set *B* = *A* \cup { $x_{i(A)+1}$ }. Let *D*(*A*) = { $x_{i(A)}, x_{h(A)}$ } (and *j*(*A*) < *h*(*A*)). By construction $D(\{x_k | k \ge i(A)\}) = \{x_{i(A)}, x_{i(A)+1}\}$ and by *A*1 *D*(*B*) = $\{x_{i(A)}, x_{i(A)+1}\}$. By *A*1 $x_{i(A)+1} \in D(B - \{x_{i(A)}\})$. Since $x_{h(A)} \in D(B - \{x_{i(A)+1}\}) = D(A)$. Let $C = B - \{x_{i(A)}\}$. By $A2 D(C) = \{x_{i(A)+1}, x_h\}$. Since $i(C) = i(A) + 1$ and $j(C) = j(A)$ we have $j(C) - i(C) < j(A) - i(A)$.

Problem A8 (Tel Aviv 2009. Inspired by Mandler, Manzini and Mariotty, (2010)).

Consider a decision maker who is choosing an alternative from subsets of a finite set X using the following procedure:

Following a fixed list of properties (a checklist), he examines one property at a time and deletes from the set all the alternatives that do not satisfy this property. When only one alternative remains, he chooses it.

1. Show that if this procedure induces a choice function, then it is consistent with the rational man model.

Let $a \in B \subset A$ such that a is the chosen alternative from A . Any other alternative in *A*, and in *B* in particular, fails to satisfy at least one of the first *m* properties that *a* does satisfy. Therefore, a is the surviving alternative also in B and condition α holds.

2. Show that any rational decision maker can be described as if he follows this procedure.

Order all the alternatives in *X* in ascending order. For each alternative $x \in X$, define the property "not being *x*" and order the properties according to the same order. Clearly, the only alternative for which "not being *x*" does not hold is *x* itself.

Let $A \subset X$ be the set from which the individual chooses. In the first stage, he deletes from *A* the worst element in *X*, if it belongs to *A*, and does nothing otherwise. Similarly, at any subsequent stage, if the alternative is not in *A* he continues to the next stage and if it does belong to *A* he deletes it. Hence, at each stage he deletes the worst alternative from his choice set. This process continues until he is left with the best alternative in *A*.

Problem A9 (Tel Aviv 2010)

A decision maker has a preference relation over \mathbb{R}^n_+ . A vector (x_1, x_2) is *interpreted as an income combination where xi is the dollar amount the decision maker receives at period i.*

Let P be the set of all preference relations satisfying:

(i) Strong Monotonicity (SM) in x_1 and x_2 .

(ii) Present preference (PP): $(x_1 + \varepsilon, x_2 - \varepsilon) \ge (x_1, x_2)$ for all $\varepsilon > 0$.

Define $(x_1, x_2)D(y_1, y_2)$ *if* $(x_1, x_2) \ge (y_1, y_2)$ for all $\ge \in P$.

1. Interpret the relation D. Is it a preference relation?

D is a domination relation: *x* dominates *y* if for every monotonic present biased preference relation, *x* is considered at least as good as *y*.

D is not a preference relation: although it is transitive, it is not complete.

2. Is it true that $(1,4)D(3,3)$? *What about* $(3,3)D(1,4)$?

 $(1,4)D(3,3)$ is false: consider the preference relation represented by the utility function $u(x_1, x_2) = x_1 + x_2$. It satisfies the two properties, but $(1, 4) \prec (3, 3)$.

 $(3,3)D(1,4)$ is true: for any preference relation which satisfies the two

properties, by PP $(3,3) \ge (1.5,4.5)$ and by SM, $(1.5,4.5) \ge (1,4)$

3. Find and prove a proposition of the following type:

 $(x_1, x_2)D(y_1, y_2)$ if and only if [put here a condition on (x_1, x_2) and (y_1, y_2)]. **Proposition**: $(x_1, x_2)D(y_1, y_2)$ Iff $x_1 \ge y_1$ and $(x_1 + x_2) \ge (y_1 + y_2)$. **Proof:**

 ξ Assume $x_1 \ge y_1$ and $(x_1 + x_2) \ge (y_1 + y_2)$.

Let $≥∈$ P, i.e. a preference relation satisfying SM and PP. If $x_2 ≥ y_2$, by SM *x* \ge *y*. If *y*₂ > *x*₂ let ϵ = *y*₂ − *x*₂.

It follows that $x_1 \ge y_1 + \epsilon$. By SM $(x_1, x_2) \ge (y_1 + \epsilon, x_2)$ and by PP $(y_1 + \epsilon, x_2) = (y_1 + \epsilon, y_2 - \epsilon) \geq (y_1, y_2).$

 \Rightarrow Assume $(x_1, x_2)D(y_1, y_2)$.

The condition $x_1 \ge y_1$ is necessary: let *a* and *b* be two vectors such that *aDb* and $a_1 < b_1$. Consider the preferences \succcurlyeq_i represented by $u(x_1, x_2) = tx_1 + x_2$ where $t > 1$. Obviously they satisfy PP and SM. For *t* large enough $ta_1 + a_2 < tb_1 + b_2$ and thus $b \succ_t a$.

The condition $x_1 + x_2 \ge y_1 + y_2$ is necessary: let *a* and *b* be two vectors such that *aDb* and $a_1 + a_2 < b_1 + b_2$. Then $b > a$.

Problem A10 (Tel Aviv 2003. Based on Gilboa and Schmeidler (1995).)

An agent must decide whether to do something, Y, or not to do it, N.

A history is a sequence of results for past events in which the agent chose Y; each result is either a success S or a failure F. For example, S,*S*,*F*,*F*,*S is a history with five events in which the action was carried out. Two of them (events* 3 *and* 4*) ended in failure while the rest were successful.*

The decision rule D is a function that assigns the decision Y or N to every possible history.

Consider the following properties of decision rules:

*A*1 : *After every history that contains only successes, the decision rule will dictate Y and after every history that contains only failures, the decision rule will dictate N.*

*A*2 : *If the decision rule dictates a certain action following some history, it will dictate the same action following any history that is derived from the first history by reordering its members. For example,* $D(S, F, S, F, S) = D(S, S, F, F, S)$.

 $A3:$ *If* $D(h) = D(h')$, then this will also be the decision following the *concatenation of h and h'. (Reminder: The concatenation of* $h = (F, S)$ *and* $h' = (S, S, F)$ *is* (F, S, S, S, F) .

1. *For every* $i = 1,2,3$, give an example of a decision rule that does not fulfill *property Ai but does fulfill the other two properties.*

A2 and A3 but not A1: Choose always *Y*. A1 is violated since $D(F) = Y$.

A1 and A3 but not A2: Choose *Y* if the first experience was *S* and choose *N* otherwise. A2 is violated since $D(S, F) = Y$ while $D(F, S) = N$.

A1 and A2 but not A3: Choose *Y* after (*S*) or if the experience does not contain two failures. A3 is violated since $D(F, S) = Y$ but $D(F, S, F, S) = N$.

2. *Give an example of a decision rule that fulfills all three properties.*

For a given $\alpha > 0$, choose *Y* iff at least α of the cases in the experience were *S*.

3. *(Difficult) Characterize the decision rules that fulfill the three properties.*

Claim: A decision rule *D* satisfies the three axioms iff there is $1 \ge \alpha \ge 0$, such that it chooses Y if the proportion of S is above α and chooses N if the proportion of S is below α .

Proof: Denote by $q(h)$ the proportion of *S* in the history *h*, let $n(h)$ be its length and let *kh* be the replication of *h*, *k* times. Assume by contradiction that there are two histories x and y such that $q(x) \geq q(y)$, $D(y) = Y$ and $D(x) = N$.

Using A3 $D(n(y)x) = N$, $D(n(x)y) = Y$ and the length of the two histories is the same $(n(x)n(y))$. Thus, we can assume that $n(x) = n(y)$.

By A2 we can also assume WLOG that all the failures appear in the sequences at the end of the sequences. Thus we get something like

 $x = (S, S, S, S, S, F, F)$

 $y = (S, S, S, F, F, F, F)$.

The number of *S*'s in *x* is larger by $k > 0$ than in *y*. Thus by A3 $D(x, kF) = N$ and $D(kS, y) = Y$ though $(x, kF) = (kS, y)$ a contradiction.

Note that for any $\alpha \in \mathbb{Q}$, the decision rule has to specify also a choice for the case when the proportion of *S* is exactly α . For $1 > \alpha > 0$, the decision rule can be either *Y* or *N*, for $\alpha = 1$ it has to be *Y* and for $\alpha = 0$ it has to be *N*. Since the proportion of *S* is a rational number, it will never be equal to α if $\alpha \notin \mathbb{Q}$.

Problem A11 (Tel Aviv 2011.)

You have read an article in a "prestigious" journal about a decision maker (DM) whose mental attitude towards elements in a finite set X is represented by a binary relation , which is a-symmetric and transitive but not necessarily complete*. The incompleteness is the result of an assumption that a DM is sometimes unable to compare between alternatives.*

Another, presumingly stronger, assumption made in the article is that the DM uses the following procedure: he has n criteria in mind, each represented by an ordering (a-symmetric, transitive and complete \rightarrow \in $(i = 1, ..., n)$ *. The DM decides that* $x > y$ *if and only if* $x > y$ *for every i.*

1.Verify that the relation generated by this procedure is a-symmetric and transitive. Try to convince a reader of the paper that this is an attractive assumption by giving a "real life" example in which it is "reasonable" to assume that a DM uses such a procedure in order to compare between alternatives.

 \ge is a-symmetric: If $x \ge y$ then by definition, $x \ge_i y$ for every *i*. Since \ge_i are a-syemmtric, $y \nless i$ *x* for all *i*, and by definition also $y \nless i$.

 $>$ is transitive: Let $x > y$ and $y > z$. By definition, $x > i$ *y* and $y > i$ *z* for every *i*. Since \geq_i are transitive, also $x \geq_i z$ for all *i*, and by definition $x \geq z$.

An example: A parent who considers destinations for a family vacation who ranks the different destinations according to the orderings of his children: he prefers A to B iff all his children prefer A to B.

It can be claimed that the additional assumption regarding the procedure that generates is not a "serious" one since given any *asymmetric and transitive relation,* \ge , one can find *a set of complete orderings* \ge ₁,..., \ge _n *such that* $x \ge y$ *iff* $x \succ_i y$ *for every i.*

2. Demonstrate this claim for the relation on the set $X = \{a, b, c\}$ according to *which only* $a > b$ *and the comparison between [b and c] and [a and c] are not determined.*

Let $a \succ_1 b \succ_1 c$ and $c \succ_2 a \succ_2 b$. The two relations agree only on $a \succ_i b$.

3. (Main part of the question) Prove this claim for the general case.

Guidance (for c): given an asymmetric and transitive relation on an arbitrary X, define a set of complete orderings $\{\succ_i\}$ and prove that $x \succ y$ iff for every *i*, $x \succ_i y$.

First, note that if *X* is a finite set and *P* is a asymmetric and transitive relation on *X* then *P* does not have any cycles and thus *P* can be extended to a complete ordering of *X* (see Problem Set 1).

Let Λ be the set of all complete orderings which extends \succ . We will see that $a > b$ if and only if $a > i$ *b* for all $\geq i \in \Lambda$:

(i) If $a > b$, then $a > i$ *b* for all *i* since any $\geq i \in \Lambda$ is an extension of \geq .

(ii) If not $a > b$, then let \succ^* be the relation \succ extended to include also $b \succ^* a$. The relation \succ^* does not have cycles: if there is a cycle $x_1 \succ^* ... \succ^* x_n = x_1$ then

(a) if for some *i* we have $x_i = b \succ^* a = x_{i+1}$ then since

 $a = x_{i+1}$ \geq^* x_{i+2} ... \geq^* $x_n = x_1$ \geq^* ... \geq^* $x_i = b$ by transitivity $a \geq b$ contradicting the assumption.

(b) otherise, by thranstivity $x_1 > x_2$ but also $x_2 > x_1$ conradicting asymettry. Thus, \succ^* can be extended to a complete ordering \succ' which will be an extension of \succ as well. Hence, there is an extension $\succ' \in \Lambda$ for which not $a \succ' b$.

B. The Consumer and The Producer

Problem B1 (Princeton 2002)

Consider a consumer with a preference relation in a world with two goods, X (an aggregated consumption good) and M ("membership in a club," for example), which can be consumed or not. In other words, the consumption of X can be any nonnegative real number, while the consumption of M must be either 0 *or* 1*. Assume that the consumer's preferences are strictly monotonic, continuous and satisfy the following property:*

Property E: For every, x there is y such that $(y,0) \succ (x,1)$ *(that is, there is always some amount of the aggregated consumption good that can compensate for the loss of membership).*

a. Show that any consumer's preference relation can be represented by a utility function of the type:

$$
u(x,m) = \begin{cases} x & \text{if } m = 0 \\ x + g(x) & \text{if } m = 1 \end{cases}.
$$

For any *x* let $h(x)$ satsfies $(x, 1) \sim (h(x), 0)$. Notice that such $h(x)$ always exists and it is unique. This is because $(0,0) \prec (x,1)$ by monotonicity and $(y,0) \succ (x,1)$ for some *y* by property E so continuity implies that $(x,1) \sim (z,0)$ for some *z*. Also, it must be unique because of monotonicity. The function *h* is increasing.

Let $g(x) = h(x) - x$. Let's verify that u actually represents \succeq . Case 1 $(x,0) \ge (x',0)$, iff $x \ge x'$ iff $u(x,0) = x \ge x' = u(x',0)$ Case 2 $(x, 1) \ge (x', 1)$ iff $x \ge x'$ iff $h(x) \ge h(x')$ iff $u(x, 1) = h(x) \ge h(x') = u(x', 1)$ Case 3 $(x, 1) \ge (\le)(x', 0)$ iff $(h(x), 0) \ge (\le)(x', 0)$ iff $h(x) \ge x'$ iff $u(x, 1) = h(x) \geq (\leq) x' = u(x', 0)$

Therefore, u represents \succeq .

b. (Less easy) Show that the consumer's preference relation can also be represented by a utility function of the type:

$$
u(x,m) = \begin{cases} f(x) & \text{if } m = 0 \\ f(x) + v & \text{if } m = 1 \end{cases}.
$$

Define $h^{0}(x) = x$ and $h^{n+1}(x) = h(h^{n}(x))$. Since $(x, 1) \sim (h(x), 0) \succ (x, 0)$, $h^{n+1}(x) > h^n(x)$ for all *n*.

(1) For $x \in [0, h(0)]$ let $f(x)$ be an arbitrary increasing function with $f(0) = 0$. Let $v = f(h(0)).$

(2) Continue inductively: assume that $f(x)$ has been already defined for $x \in [0, h^{n-1}(0)]$ such that $f(x)$ is increasing in the region $[h(0), h^{n-1}(0)]$ and $f(h^k(0)) = kv$ for all $k \leq n-1$. Define $f(x)$ for $x \in (h^{n-1}(0), h^n(0))$, such that $f(x) = v + f(h^{-1}(x))$. Since *h* is an increasing function, h^{-1} exists, increasing and

 $h^{-1}(x) \in (0, h(0))$. The function *f* is increasing in $x \in [h^{n-1}(0), h^{n}(0)]$ and $f(h^n(0)) = nv.$

(3) Let $u(x, 0) = f(x)$ and $u(x, 1) = f(x) + v$.

We have to make sure that *f* is actually defined for all $s \geq 0$. Since $\{h^n(0)\}\)$ is an increasing sequence, $f(x)$ is not defined twice or more in the above steps. Therefore, it is sufficient to show that $\lim_{n\to\infty} h^n(0) = \infty$.

Suppose this is not, since $\{h^n(0)\}\$ is an increasing sequence, let $sup_n h^n(0) = K$ and $h^{n}(0) < K$ for all *n*. This means that $(h^{n}(0), 1) \sim (h^{n+1}(0), 0) \prec (K, 0)$ for all *n* where the first indifference comes from the definition of *h* and the second preference comes from monotonicity. By continuity and since $h^n(0) \rightarrow K$, $(K,1) \preceq (K,0)$, contradicting monotonicity.

Finally, we need to confirm that this u actually represents \geq . Notice that u represents \succsim correctly between $(x,0)$ and $(x',0)$ (or $(x,1)$ and $(x',1)$) because *f* is an increasing function. Therefore, we need to show that $(x,1) \geq (x',0)$ if and only if $f(x) + v \ge f(x')$.

Suppose that $(x,1) \geq (x',0)$. Then, by the definition of *h*, $h(x) \geq x'$. Since *f* is increasing, we have $u(h(x),0) = f(h(x)) \ge f(x') = u(x',0)$. By construction of *f*, we $\text{have } f(h(x)) = v + f(h^{-1}(h(x))) = v + f(x) = u(x, 1).$ Therefore, $u(x, 1) \ge u(x', 0).$ The same argument can be applied for the case when $(x, 1) \precsim (x', 0)$. Hence, we conclude that *u* defined above actually represents \geq .

c. Explain why continuity and strong monotonicity (without prperty E) are not sufficient for (a)

Consider the lexicographic preference \succsim which gives a priority to M. (so (x,m) \ge (x',m') if and only if $m > m'$ or " $m = m'$ and $x > x''$ ").

This is clearly strictly monotonic both in *x* and *m*.

Continuity: To see this, take any $(x, m) \succ (x', m')$.

Suppose $m = 1$ and $m' = 0$. Take $\epsilon = 1/2$, then $(a, n) \in B_{\epsilon}((x, m))$ implies $n = 1$ and $(a', n') \in B_\epsilon((x', m'))$ implies $n' = 0$. Therefore, $(a, n) > (a', n')$. WHen $m = m'$ (so $f(x, x, y')$ let $\epsilon = (x - x')/3$. Then, $(a, n) \in B_{\epsilon}(x, m)$ and $(a', n') \in B_{\epsilon}((x', m'))$ imply that $n = n' = m$ but $a > a'$, so we have $(a, n) > (a', n')$.

However, \succeq cannot be represented by a utility function with the form given in (A) since $(0,1)$ would be indifferent to $(g(0),0)$.

d. Calculate the consumer's demand function.

For simplicity assume $p_x = 1$. By monotonicity, the consumer always spends all wealth on *X* or *M*. If $p_m \leq w$, his choice is between "buying *M* and spending all the remaining wealth on X" and "Spending all wealth on X". If $p_m > w$, he has no choice except "spending all the wealth on *X*". Therefore, his demand function is characterized by

$$
(x(p,w), m(p,w)) = \begin{cases} (w,0) & \text{if } (w,0) \ge ((w-p_m)/p_x,1) \text{ or } p_m > w \\ (w-p_m,1) & \text{if } (w,0) \le ((w-p_m)/p_x,1) \text{ and } p_m \le w \end{cases}.
$$

e. Taking the utility function to be of the form described in (1), derive the consumer's indirect utility function. For the case where the function *g* **is differentiable, verify Roy's identity with respect to commodity** *M***.**

If the consumer's utility function is given by a differentiable utility function as in part (A), then his/her indirect utility function is

$$
v(p, w) = \begin{cases} w & \text{if } w \ge (w - p_m) + g((w - p_m)) \text{ or } p_m > w \\ (w - p_m) + g((w - p_m)) & \text{if } (w, 0) \le ((w - p_m), 1) \text{ and } p_m \le w \end{cases}
$$

In the first case,

$$
\frac{\partial v/\partial p_m}{\partial v/\partial w} = 0 = m(p, w)
$$

and in the second case,

$$
\frac{\partial v}{\partial p_m} = -\frac{(-1) - g'(w - p_m)}{1 + g'(w - p_m)} = 1 = m(p, w)
$$

so the Roy equality holds.

Problem B2 (Princeton 2001)

(a) Define a formal concept for " \lesssim_1 *and* \gtrsim_0 *are closer than* \gtrsim_2 *and* \gtrsim_0 ".

Let us say that \gtrsim_1 and \gtrsim_0 are closer than \gtrsim_2 and \gtrsim_0 *y* and $x \gtrsim_0 y$ imply $x \geq 1$ *y*, and $x \geq 2$ *y* and $x \geq 0$ *y* imply $x \geq 1$ *y*.

(b) Apply your definition to the class of preference relations represented by $U_1 = tU_2 + (1-t)U_0$, where the function U_i represents \succeq_i $(i = 0,1,2)$.

Let $0 \le t \le 1$. If $U_2(x) \ge \lceil 5 \rceil U_2(y)$ and $U_0(x) \ge \lceil 5 \rceil U_0(y)$, then $U_1(x) = tU_2(x) + (1-t)U_0(x) \geq [t]U_2(y) + (1-t)U_0(y) = U_1(y)$. So the preferences represented by U_1 and the preferences represented by U_0 are closer than the preferences represented by U_2 and the preferences represented by U_0 .

(c) Consider the above definition in the consumer context. Denote by $x_k^i(p, w)$ *the demand function of* ζ_i *for good k. Show that* ζ_1 *and* ζ_0 *may be closer than* \succsim_2 and \succsim_0 , and nevertheless $|x_k^1(p,w)-x_k^0(p,w)|> |x_k^2(p,w)-x_k^0(p,w)|$ for some *commodity k, price vector p and wealth level w*.

Let $K = 3$, $U_0(x) = \min\{x_1, x_2\}$ and $U_2(x) = \min\{x_2, x_3\}$ and let U_1 be defined as in (b) with $t = 1/2$. Then for $p = (1,1,1)$ and $w = 1$, the demands associated with U_0 , *U*₁ and *U*₂ are $x_0 = (1/2, 1/2, 0), x_1 = (1/3, 1/3, 1/3)$ and $x_2 = (0, 1/2, 1/2)$ respectively. By (b), the preferences induced by U_1 and the preferences induced by U_0 are colser than the preferences induced by U_2 and the preferences induced by U_0 , but $|x_2^1 - x_2^0| = 1/6 > 0 = |x_2^2 - x_2^0|.$

Problem B3 (Tel Aviv 2003)

Consider the following consumer problem: There are two goods, 1 and 2. The consumer has a certain endowment. His preferences satisfy monotonicity and continuity. Before the consumer are two "exchange functions": he can exchange x units of good 1 for $f(x)$ *<i>units of good 2, or he can exchange* y *units of good 2 for gy units of good 1. Assume the consumer can only make one exchange.*

1. *Show that if the exchange functions are continuous, then a solution to the consumer problem exists.*

Denote by *w* the initial endowment. the budget set includes $\{(x_1, x_2) | x_1 \leq w_1 \}$ and $(x_2 - w_2) = f(w_1 - x_1)$, or, $x_1 \geq w_1$ and $(x_1 - w_1) = g(w_2 - x_2)$. This is a compact set and thus the consumer's preferences has a maximum point.

2. *Explain why strong convexity of the preference relation is not sufficient to guarantee a unique solution if the functions f and g are increasing and convex.*

Take, for example, $w = (4, 4)$, $f(x) = 2x$ and $g(y) = 2y$. Consider the utility function $u(x_1, x_2) = x_1x_2$. Clearly both $(6, 3)$ and $(3, 6)$ are solutions.

3. *Interpret the statement "the function f is increasing and convex".*

The more units the consumer will exchange of commodity 1 the better will be is exchange rate.

4. *Suppose both functions f and g are differentiable and concave and that the product of their derivatives at point* 0 *is* 1*. Suppose also that the preference relation is strongly convex. Show that under these conditions, the agent will not find two different exchanges, one exchanging good 1 for good 2, and one exchanging good 2 for good 1, optimal.*

Since $f'(0) = 1/g'(0)$, the consumer's budget line is differentiable at (w_1, w_2) , which is his original endowment (note that *f* is a function from good 1 to good 2 while *g* is a function from good 2 to good 1). The functions *f* and *g* are both concave, thus the budget line is convex. Assume, in contradiction, that the consumer finds two exchanges to be optimal, one that yields the consumption bundle (x_1, x_2) and one that yields (y_1, y_2) . Since the budget line is convex, the bundle ($\frac{x_1+y_1}{2}, \frac{x_2+y_2}{2}$) is in the consumer's budget set. Since the consumer is indifferent between (x_1, x_2) and (y_1, y_2) , by the strict convexity of his preference relation he would strictly prefer the bundle $(\frac{x_1+y_1}{2}, \frac{x_2+y_2}{2})$, in contradiction to (x_1, x_2) and (y_1, y_2) being optimal.

If it would be the case it would mean that there are two exchanges $(-a, +f(a))$ and $(Fg(b), -b)$ which are better than $(0, 0)$. By the concavity of *f* and *g*, $f(a) < a$ and $g(b) < b$. Let $\lambda/(1 - \lambda) = b/a$. By convexity the change $(-\lambda a + (1 - \lambda)g(b), +\lambda f(a) - (1 - \lambda)b)$ is improving but $-\lambda a + (1 - \lambda)g(b) < -\lambda a + (1 - \lambda)b = 0$ and $\lambda f(a) - (1 - \lambda)b < \lambda a - (1 - \lambda)b = 0$ a contradiction to the monotonicity of the preferences.

5. *Now assume* $f(x) = ax$ and $g(y) = by$ *. Explain this assumption. Find a condition that will ensure it is not profitable for the consumer to make more than one exchange.*

If the consumer sells one unit of the first commodity he gets *a* of the second and exchanging it back to commodity 2 he will get *ab* units of commodity 1. It is

necessary that $ab < 1$. Of course if $ab > 1$ he would be able to reach any bundle.

Problem B4 (Tel Aviv 1998)

A consumer with wealth w 10 *"must" obtain a book from one of three stores. Denote the prices at each store as* p_1, p_2, p_3 . All prices are below w in the relevant *range. The consumer has devised a strategy: he compares the prices at the first two stores and purchases the book from the first store if its price is not greater than the price at the second store. If* $p_1 > p_2$, he compares the prices of the second and *third stores and purchases the book from the second store if its price is not greater than the price at the third store. He uses the remainder of his wealth to purchase other goods.*

1. *What is this consumer's "demand function"?*

Denote a bundle by (m, x_1, x_2, x_3) where *m* is remaining wealth, and x_i is 1 if he purchases the book from the *i*'th store and 0 otherwise.

$$
(w - p_1, 1, 0, 0) \quad \text{if } p_1 \le p_2
$$
\n
$$
\text{The demand function } x(w, p_1, p_2, p_3) \text{ is } (w - p_2, 0, 1, 0) \quad \text{if } p_1 > p_2 \le p_3
$$
\n
$$
(w - p_3, 0, 0, 1) \quad \text{if } p_1 > p_2 > p_3
$$

2. *Does this consumer satisfy "rational man" assumptions?*

No. Under the price vector $(3,3,1)$, the consumer buys the book in the first shop and his consumption bundle is $(w - 3, 1, 0, 0)$ whereas $(w - 1, 0, 0, 1)$ was feasible.

Under the price vector $(3,2,1)$, the consumer buys the book in the third shop, the bundle $(w - 1, 0, 0, 1)$ whereas $(w - 3, 1, 0, 0)$ was feasible.

3. *Consider the function* $v(p_1, p_2, p_3) = w - p_i^*$, where i^* is the store from which *the consumer purchases the book if the prices are* (p_1, p_2, p_3) . What does this *function represent?*

Since the consumer must buy the book, his welfare is measured by the money left in his pocket. Thus, *v* could be thought of as an indirect utility function.

4. *Explain why v is not monotonically decreasing in pi. Compare with the indirect utility function of the classic consumer model.*

If $p_3 < p_1 < p_2$ then an increase in p_1 will improve the consumer's welfare. In contrast, an indirect utility function which is derived from a maximization of a utility function is always non-increasing in a price.

Problem B5 (NYU 2005)

A consumer has preferences which satisfy monotonicity, continuity and strict convexity, in a world of K goods. The goods are split into two categories, 1 *and* 2*, of* K_1 *and* K_2 *goods respectively* $(K_1 + K_2 = K)$ *. The consumer receives two types of money: wi units of money of type i*, *which can only be exchanged for goods in the i'th category given a price vector* p_i *.*

Define the induced preference relation over the two-dimensional space (w_1, w_2) . *Show that these preferences are monotonic, continuous and convex.*

The consumer's problem is to mximize the preferences \geq over the set $\{(x_1, x_2) |$ $x_1 \in \mathbb{R}^{K_1}, x_2 \in \mathbb{R}^{K_2}, p_1x_1 \leq w_1$ and $p_2x_2 \leq w_2$. Denote its solution by $x(w_1, w_2)$. Define the preference relation over the space (w_1, w_2) as

$$
(w_1,w_2) \succcurlyeq^* (w_1',w_2') \Leftrightarrow x(w_1,w_2) \succcurlyeq x(w_1',w_2')
$$

Now let's prove the properties of \succcurlyeq^* .

Monotonicity: Let $(w'_1, w'_2) \gg (w_1, w_2)$. This implies that

$$
p_1x_1(w_1,w_2) < w_1' \text{ and } p_2x_2(w_1,w_2) < w_2'.
$$

The budget conatraints contain a bundle *y* which "dominates" $x(w_1, w_2)$. Therefore, by Monotonicity of \succsim we get $x(w'_1,w'_2) > x(w_1,w_2)$, that is $(w'_1,w'_2) >^* (w_1,w_2)$.

Convexity: Let $(w'_1, w'_2) \geq^* (w_1, w_2)$. This implies that $x(w'_1, w'_2) \geq x(w_1, w_2)$. Let $w^{\lambda} = \lambda(w_1', w_2') + (1 - \lambda)(w_1, w_2)$ and $x^{\lambda} = \lambda x(w_1', w_2') + (1 - \lambda)x(w_1, w_2)$ for some $\lambda \in [0,1]$. We know that

$$
p_1x_1(w_1, w_2) \le w_1 \text{ and } p_2x_2(w_1, w_2) \le w_2
$$

$$
p_1x_1(w'_1, w'_2) \le w'_1 \text{ and } p_2x_2(w'_1, w'_2) \le w'_2
$$

which implies that

$$
p_1x_1^{\lambda} \leq w_1^{\lambda}, p_2x_2^{\lambda} \leq w_2^{\lambda}.
$$

Moreover, convexity of \geq implies that $x^{\lambda} \geq x(w_1, w_2)$. Therefore, it must be that $x(w_1^{\lambda}, w_2^{\lambda}) \succsim x^{\lambda} \succsim x(w_1, w_2)$, which implies that $(w_1^{\lambda}, w_2^{\lambda}) \succsim^* (w_1, w_2)$.

Continuity: To show this we'll first show that the demand function *x* is continuous. So, let $(w_1^n, w_2^n) \rightarrow (w_1, w_2)$. We need to show that $x(w_1^n, w_2^n) \rightarrow x(w_1, w_2)$. Note first that the sequence $x(w_1^n, w_2^n)$ is inside of some compact set: define

$$
\overline{w}_1 = \sup \{w_1^n\}, \overline{w}_2 = \sup \{w_2^n\}.
$$

It's clear that for any *n* we must have

$$
x(w_1^n, w_2^n) \in \{x \in \Re_+^{k_1+k_2} : p_1x_1 \leq \overline{w}_1 \text{and } p_2w_2 \leq \overline{w}_2\}.
$$

So our sequence is indeed entirely contained in a compact set. Now suppose that $x(w_1^n, w_2^n) \nrightarrow x(w_1, w_2).$ This implies that there exists a subsequence $x(w_1^k, w_2^k),$ such that $x(w_1^k, w_2^k) \rightarrow y \neq x(w_1, w_2)$. The fact that $x(w_1^k, w_2^k) \rightarrow y$ implies that

$$
p_1y_1 \leq w_1, p_2y_2 \leq w_2.
$$

By the strict convexity, $x(w_1, w_2) \succ (y_1, y_2)$. Continuity of \succcurlyeq implies that there exists $z \ll x(w_1, w_2)$ such that $z \succ (y_1, y_2)$. But then, continuity of \geq together with the fact that $p_1z_1 < w_1$ and $p_2z_2 < w_2$ imply that for *k* large enough

$$
p_{1}z_{1} < w_{1}^{k}, p_{2}z_{2} < w_{2}^{k}, z > x(w_{1}^{k}, w_{2}^{k})
$$

which contradicts the optimality of $x(w_1^k, w_2^k)$.

Now that we know that the demand function is continuous we can easily show that \succ * is continuous. Let $(w_1^n, w_2^n) \to (w_1, w_2)$ and $(\hat{w}_1^n, \hat{w}_2^n) \to (\hat{w}_1, \hat{w}_2)$ be such that $(w_1^n, w_2^n) \geq^* (\hat{w}_1^n, \hat{w}_2^n)$ for all *n*. This implies that $x(w_1^n, w_2^n) \geq x(\hat{w}_1^n, \hat{w}_2^n)$ for all *n*. By continuity of \succcurlyeq and continuity of the demand function *x*, $x(w_1, w_2) \succeq x(\hat{w_1}, \hat{w_2})$, which is equivalent to $(w_1, w_2) \geq^* (\hat{w_1}, \hat{w_2})$.

Problem B6 (NYU 2006)

Consider a consumer in a world of 2 commodities who has to make choices from budget sets parametrized by (p, w) , with the additional constraint that the *consumption of good 1 is limited by some external bound* $c \geq 0$. That is, in his *world, a choice problem is a set of the form* $B(p, w, c) = \{x|px \leq w \text{ and } x_1 \leq c\}$ *. Denote by* $x(p, w, c)$ *the consumer's choice from* $B(p, w, c)$ *.*

1. Assume that $px(p, w, c) = w$ and $x_1(p, w, c) = \min\{0.5w/p_1, c\}$. Show that this *behavior is consistent with the assumption that demand is derived from a maximization of some preference relation.*

Maximization the preferences \succsim represented by the Cobb-Douglas utility function $u(x) = x_1^{0.5}x_2^{0.5}$ leads to the given demand function $x(p, w, c)$. There are two different cases to consider.

Case 1: 0.5*w*/*p*¹ ≤ *c*

In this case, $x(p, w, c)$ is equal to the point $(0.5w/p₁, 0.5w/p₂)$, which is the unique maximizer of *u* on the usual budget set $B(p, w) = \{x \mid px \leq w\}$. In particular, $x(p, w, c)$ is the unique maximizer of *u* on $B(p, w, c)$.

Case 2: $0.5w/p_1 > c$

Suppose by contradiction that $x(p, w, c) = (c, (w - p_1 c)/p_2)$ is not the unique maximizer of *u* on $B(p, w, c)$. Then there is a $y \in B(p, w, c)$ with $y \neq x(p, w, c)$ such that $y \geq x(p, w, c)$. By strict monotonicity of *u* we must have $y_1 < c$, otherwise we would have $x(p, w, c) > v$. Moreover, we can assume $pv = w$. Define $\bar{x} = (0.5w/p_1, 0.5w/p_2)$. Now, since $y_1 < c < 0.5w/p_1$, $x(p,w,c)$ can be written as a strict convex combination of the points *y* and \bar{x} . Since \bar{x} is the unique maximizer of *u* on $B(p, w)$, by strict convexity of \succeq we must have $x(p, w, c) > y$, a contradiction.

2. Assume that $px(p, w, c) = w$ and $x_1(p, w, c) = \min\{0.5c, w/p_1\}$. Show that this *consumer's behavior is inconsistent with preference maximization.*

Suppose by contradiction that $x(p, w, c)$ is consistent with maximization of a preference relation \geq . Fix a price vector p and wealth level w. Pick a $c < w/p_1$, so that $x(p, w, c) = (0.5c, (w - p_1 0.5c)/p_2)$. Now we can pick a c' , sufficiently close to c , such that

$$
0.5c' < c < c' < w/p_1.
$$

Since $0.5c' < w/p_1$, $x(p,w,c') = (0.5c', (w-p_1 0.5c')/p_2) \neq x(p,w,c)$. Moreover, since $0.5c' < c$, we have $x(p, w, c') \in B(p, w, c)$, and hence $x(p, w, c') \prec x(p, w, c)$, a contradiction to the optimality of $x(p, w, c')$.

3. *Assume that the consumer chooses his demand for x by maximizing the utility function* $u(x)$ *. Denote the indirect utility by* $V(p, w, c) = u(x(p, w, c))$ *. Assume V is "well-behaved". Outline the idea of how one can derive the demand function from the function V in case that* $\partial V/\partial c(p, w, c) > 0$.

Fix a parameter vector $t^* = (p^*, w^*, c^*)$ and assume $\partial V(t^*)/\partial c > 0$. We claim that $x_1(t^*) = c^*$.

Suppose by contradiction that $x_1(t^*) < c^*$. However, since $\partial V(t^*)/\partial c > 0$, there

exists an $\epsilon > 0$ such that $V(p^*, w^*, c^* - \epsilon) < V(t^*)$ and $c^* - \epsilon > x_1(t^*)$. Since $p^*x(t^*) = w^*$, it follows that $x(t^*) \in B(p^*, w^*, c^* - \epsilon)$. Hence, $V(t^*) = u(x(t^*)) \leq V(p^*, w^*, c^* - \epsilon)$, a contradiction to $\partial V(t^*)/\partial c > 0$.

Problem B7 (Tel Aviv 2006)

Imagine a consumer who lives in a world with $K + 1$ *commodities and behaves in the following manner: The consumer is characterized by a vector D, consisting of the commodities* 1,..,*K. If he can purchase D, he will consume it and spend the rest of his income on commodity* $K + 1$. If he is unable to purchase D, he will not *consume commodity* $K + 1$ *and will purchase the bundle tD* $(t \leq 1)$ where *t* is as *large as he can afford.*

1. *Show that there exists a monotonic and convex preference relation which explains this pattern of behavior.*

Consider the preferences represnted by

$$
u(x) = \begin{cases} 1 + x_{K+1} & \text{if } x \ge D \\ \min\{x_1/d_1, \dots, x_K/d_K\} & \text{otherwise} \end{cases}
$$

Since $\min\{x_1/d_1,\ldots,x_K/d_K\} \leq 1$ for all bundles in the second range, the consumer will prefer to be in the first range so whenever he can purchase *D* he will choose a bunlde in the first range where it is optimal for him to purchase as much as possible on the $K + 1$ 'th good. If he cannot afford D then his behvaior will be as a amximizer of $\min\{x_1/d_1,\ldots,x_K/d_K\}$ and thus he will consume a bundle *tD* with the highest *t* he can afford.

2. *Show that there is no monotonic and continuous preference relation that explains this pattern of behavior.*

If there was such a preference relation it would have to satsify $(d_1,...,d_K2)$ > $(d_1+1,...,d_K+1,0)$. However, $(d_1,...,d_K0)$ > $(d_1-\delta,d_2,...,d_K2)$ for all $\delta > 0$, thus by continuity $(d_1,...,d_K,0) \geq (d_1,...,d_K,2)$ contradicting monotonicity.

Problem B8 (NYU 2007)

A consumer in a world of K commodities maximizes the utility function $u(x) = \sum_{k} x_{k}^{2}$.

1. *Calculate the consumer's demand function (whenever it is uniquely defined).* Demand is uniquely defined when there is a unique minimal price p_k . The consumer will set $x_k(p, w) = \frac{w}{p_k}$ and $x_j(p, w) = 0$ for all $j \neq k$.

2. *Give another preference relation (not just a monotonic transformation of u) which induces the same demand function.*

Preferences represented by $v(x) = \sum_{k} x_k$.

3. For the original utility function u , calculate the indirect preferences for $K = 2$. *What is the relationship between the indirect preferences and the demand function?* (It is sufficient to answer for the domain where $p_1 < p_2$.)

The indirect preferences are represented by $v(p,w) = [\frac{w}{\min\{p_1,p_2\}}]^2$.

When $p_1 < p_2$, demand can be induced from v as follows:

$$
x(p,w) = \left(-\frac{\partial v(p,w)/\partial p_1}{\partial v(p,w)/\partial w}, -\frac{\partial v(p,w)/\partial p_2}{\partial v(p,w)/\partial w}\right) = \left(-\frac{-2w/p_1^2}{2/p_1}, 0\right) = \left(\frac{w}{p_1}, 0\right).
$$

4. *Are the preferences in (1) differentiable (according to the definition given in class)?*

No. Let $K = 2$ and consider the bundle $(1,1)$. The vectors $d = (1,-1)$ and $d' = (-1,1)$ are both improvement directions, but for any vector of values *v*, if $v \cdot d > 0$, then $v \cdot d' < 0$.

Problem B9 (NYU 2008)

A decision maker has a preference relation over the pairs (x_{me}, x_{him}) with the *interpretation that xme is an amount of money he will get and xhim is the amount of money another person will get. Assume that*

(i) for all (a, b) such that $a > b$ the decision maker strictly prefers (a, b) over *b*,*a.*

(ii) if $a' > a$ then $(a', b) > (a, b)$.

The decision maker has to allocate M between him and another person.

1. *Show that these assumptions guarantee that he will never allocate to the other person more than he gives to himself.*

Let $B(M) = \{(a, b) | a + b \leq M\}$ be the set of feasible allocations, and $x(M) = (x_1, x_2)$ be the chosen allocation from the feasible set, that is $x(M) \geq (a, b)$ for any $(a,b) \in B(M)$. For any $a < b$ by (i) $(b,a) > (a,b)$, and (b,a) is feasible so that (a, b) is not optimal. Therefore $x_1 \geq x_2$.

2. *Assume (i), (ii) and*

(iii) The decision maker is indifferent between (a, a) and $(a - \varepsilon, a + 4\varepsilon)$ for all a *and* $\varepsilon > 0$ *.*

Show that nevertheless he might allocate the money equally.

Suppose the preferences are represented by, for example

$$
u(x,y) = \begin{cases} 4x + y & \text{if } y \ge x \\ 2x + 3y & \text{if } x \ge y \end{cases}
$$

These preferences satisfy monotonicity.

The preferences satisfy (iii) since ∀*a*,

 $u(a,a) = 5a = 4(a-\varepsilon) + (a+4\varepsilon) = u(a-\varepsilon, a+4\varepsilon).$

They also satisfy (i) since if $x_2 > x_1$, then

 $u(x_1, x_2) = 4x_1 + x_2 = 5x_1 + (x_2 - x_1) < 5x_1 + 2(x_2 - x_1) = 2x_2 + 3x_1 = u(x_2, x_1).$

In this case, the utility would be maximized by setting $x = y = \frac{M}{2}$.

3. *Assume (i), (ii), (iii) and*

(iv) The decision maker's preferences are also differentiable (according to the definition given in class).

Show that in this case, he will allocate to himself (strictly) more than to the other.

Assume by contradiction the DM chooses to allocate $x(M) = (\frac{M}{2}, \frac{M}{2})$. By differentiability of the preferences, $v(x) = (4,1)$ (i.e. the hyperplane separating the improving directions would be the hyperplane $4x + y = M$). But then $(1, -1)$ is a strictly improving direction, and the bundle ($\frac{M}{2} + \epsilon$, $\frac{M}{2} - \epsilon$) > ($\frac{M}{2}$, $\frac{M}{2}$) = $x(M)$ would be affordable for small $\epsilon > 0$, a contradiction.

Problem B10 (NYU 2009)

An economic agent is both a producer and a consumer. He has a_0 units of good 1. He can use some of a_0 to produce commodity 2. His production function f *satisfies monotonicity, continuity, strict concavity. His preferences satisfy monotonicity, continuity and convexity. Given he uses a units of commodity* 1 *in production he is able to consume the bundle* $(a_0 - a, f(a))$ *for* $a \le a_0$ *. The agent has in his "mind" three "centers":*

**The pricing center declares a price vector* (p_1, p_2) .

**The production center takes the price vector as given and operates according to one of the following two rules*

Rule 1: maximizing profits, $p_2f(a) - p_1a$ *.*

Rule 2: maximizing production subject to the constraint of not making any losses,, i.e. $p_2f(a) - p_1a \ge 0$ *.*

The output of the production center is a consumption bundle.

The consumption center takes $(a_0 - a, f(a))$ as endowment, and finds the optimal *consumption allocation that it can afford according to the prices declared by the pricing center.*

The prices declared by the pricing center are chosen to create harmony between the other two centers in the sense that the consumption center finds the outcome of the production center's activity, $(a_0 - a, f(a))$, *optimal given the announced prices.*

1. Show that under Rule 1, the economic agent consumes the bundle $(a_0 - a^*, f(a^*))$ *which maximizes his preferences.*

The solution corresponds to the point on the production possibility set where preferences are maximized.

Since the production possibility set is strictly convex, and preferences are convex we know that there is a unique maximum.

Now choose a price vector such that the price line is tangent to this set and the indifference line exactly at the maximum.

By construction, profit is maximized given prices, and preferences are
maximized at the intersection point for given prices and endowment point $(a_0 - a^*, f(a^*)).$

2. What is the economic agent's consumption with Rule 2?

The economic agent chooses $(a_0 - a^*, f(a^*))$ with maximal a^* subject to the constraint that preferences are maximized at this point when we take the line connecting this point to $(a_0, 0)$ as the price line. By construction, production is maximized here subject to the constraint that there are no losses with the given prices. Also the point is chosen to guarantee that the consumer preferences are maximized at the budget set with the same prices.

3. State and prove a general conclusion about the comparison between the behavior of two individuals, one whose production center operates with Rule 1 and one whose production center activates Rule 2.

Claim: Individual using Rule 2 will always produce more, i.e. for a_1, p^1 and a_2, p^2 denoting the solutions under Rule 1 and Rule 2, $f(a_1) \le f(a_2)$.

Assume for contradiction that $a_1 > a_2$. This means that the solution with Rule 2 is strictly to the right of the solution with Rule 1. Since *f* is strictly concave and monotonic, if solution with Rule 2 is to the right of solution with Rule 1, we must have $\frac{p_1^1}{1}$ $\frac{p_1^1}{p_2^1} < \frac{p_1^2}{p_2^2}$ $\frac{p_{\bar{1}}}{p_2^2}$.

Note that $(a_0 - a_1, f(a_1))$ affordable (and strictly interior) in the budget set defined by p^2 . Also $(a_0 - a_2, f(a_2))$ is affordable (and strictly interior) in the budget set defined by $p¹$.

 \Rightarrow $(a_0 - a_2, f(a_2)) \rightarrow (a_0 - a_1, f(a_1))$ \Rightarrow $(a_0 - a_1, f(a_1)) \rightarrow (a_0 - a_2, f(a_2))$ which is a contradiction.

Problem B11 (Tel Aviv 2010)

A basketball coach considers buying players from a set A. Given a budget w and a price vector $(p_a)_{a \in A}$ *the coach can purchase any set such that the total cost of the players in it is not greater than w. Discuss the rationality of each of the following choice procedures, defined for any budget level w and price vector P:*

(P1) The consumer has in mind a fixed list of the players in $A: a_1, \ldots, a_n$ *.* Starting at the beginning of the list, when he arrives to the *i's* player he adds him to *the team if his budget allows him to after his past decisions, and then continues to the next player on the list with his remaining budget. This continues until he runs out of budget or has gone through the entire list.*

Let $\{1,...,n\}$ be the list of players. Identify a set *B* with a vector $x(B)$ of 0's and 1's such that $x(B)_i = 1$ if $i \in B$ and $x(B)_i = 0$ if $i \notin B$. The consumer's choice is rationalized by the preferences: $B \ge C$ if $x(B) \ge_L x(C)$, where \ge_L are the standard lexicographic preferences on \mathbb{R}^n .

(P2) He purchases the combination of players that minimize the excess budget he is left with.

The procedure is NOT rationalizable since it is even does not induce a choice from a choice set:

The choice set is $\{\{a_1\}, \{a_2\}\}\$ for both sets of parameters:

 $p_1 = 2$, $p_2 = 3$ and $w = 3$ and $p_1 = 3$, $p_2 = 2$ and $w = 3$.

But in one case $\{a_1\}$ is chosen and in the other $\{a_2\}$.

Problem B12 (Tel Aviv 2011)

A consumer in a two commodity world operates in the following manner: The consumer has a preference relation $\zeta_{\rm S}$ on \mathbb{R}^2_+ . His father has a preference *relation* \geq_F *on the space of his son's consumption bundles. Both relations satisfy strong monotonicity, continuity and strict convexity. The father does not allow his son to purchase a bundle which is not as good (from his perspective) as the bundle M*,0*. The son, when choosing from a budget set, maximizes his own preferences subject to the constraint imposed by his father. In the case that he cannot satisfy his father's wishes, he feels free to maximize his own preferences.*

1. Prove that the behavior of the son is rationalizable.

Define > as follows: $a > b$ iff (i) $a \succeq_F (M,0)$ and $(M,0) >_F b$, or (ii) both $a, b \geq_F (M, 0)$ and $a \geq_S b$ or (iii) both $(M, 0) \geq_F a, b$ and $a \geq_S b$.

 \geq can easily be shown to be complete and transitive.

1. Prove that the preferences which rationalize this kind of behavior are monotonic.

Take any *x*, *y* st. $x_1 \geq y_1$ and $x_2 \geq y_2$. Since \succsim_S , \succsim_F are monotonic, $x \succsim_S y$ and $x \succeq_F y$. Thus by construction of $\succ, x \succeq y$.

1. Show that the preferences which rationalize this kind of behavior are not necessarily continuous nor convex (you can demonstrate this diagrammatically).

To see possible violation of continuity and convexity consider the example below.

Note that $(M,0) > (0,2)$ since $(0,2)$ is below the father's indifference curve passing through *M*,0. However we can see from the son's preferences that for any $\alpha \in (0,1), (0,2) \succ \alpha(0,2) + (1-\alpha)(M,0)$ violating convexity and continuity.

4. (Bonus) Assume that the father's instructions are that given the budget set (p, w) *the son is not to purchase any bundle which is* \geq_F *-worse than* $(w/p_1, 0)$ *. The son seeks to maximize his preferences subject to satisfying his father's wishes. Show that the son's behavior satisfies the Weak Axiom of Revealed Preferences.*

Assume there is a violation of the WARP. Then there must be two overlapping budget sets as shown above such that *a* is chosen from set *A* and *b* is chosen from set *B*.

It must be that $a \succeq_F (x_2, 0)$ and $b \succeq_F (x_1, 0)$. By monotonicity, $(x_2, 0) \succeq_F (x_1, 0)$ and thus $a \succeq_F (x_1, 0)$ Since *b* is chosen over *a* in set *B*, $b \succ_S a$ By monotonicity, there exists $z \in B$ st $z \succeq_S b \succ_S a$ Also by convexity $z \succeq_F x_2$, contracting a being optimal in set *A*.

C. Uncertainty:

Problem C1 (Princeton 2001)

A consumer has to make a choice of a bundle before he is informed whether a certain event, which is expected with probability α and affects his welfare, has *happened or not. He assigns a vNM utility* $v(x)$ to the consumption of the bundle x when the event occurs, and a vNM utility $v'(x)$ to the consumption of x should the *event not occur. Having to choose a bundle the consumer maximizes his expected utility* $a v(x) + (1 - a) v'(x)$. Both *v* and v' induce preferences on the set of bundles *satisfying the standard assumptions about the consumer. Assume also that v and v*′ *are concave.*

(a)*Show that the consumer's preference relation is convex.*

The function $a v(x) + (1 - a) v'(x)$ is a convex combination of concave functions, thus it is quasi-concave and induces a preference relation which is convex.

(b) *Find a connection between the consumer's indirect utility function and the indirect utility functions derived from v and v*′ *.*

Let $x(p, w)$ denote the demand and f, f_v and $f_{v'}$ the indirect utility functions. Then, $f(p, w) = \alpha v(x(p, w)) + (1 - \alpha)v'(x(p, w)) \leq \alpha f_v(p, w) + (1 - \alpha)f_{v'}(p, w)$. The interpretation of this property is that the decision maker is better off if he makes his choice after rather than before the uncertainty is resoleved.

(c) *A new commodity appears on the market: "A discrete piece of information that tells the consumer whether or not the event occurred". The commodity can be purchased prior to the consumption decision. Use the indirect utility functions to characterize the demand function for the new commodity.*

Suppose that *v* and v' are continuous and monotonic so that *f*, f_v and $f_{v'}$ are strictly increasing and continuous in w . Then for given (p, w) , by (b), there is a unique $\beta^*(p, w)$ that satisfies:

 $f(p, w) = \alpha f_v(p, w - \beta^*(p, w)) + (1 - \alpha) f_v'(p, w - \beta^*(p, w)).$

The demand for this good is 1 if its price is below $\beta^*(p, w)$ and 0 otherwise.

Problem C2 (Tel Aviv 1999)

Tversky and Kahneman (1986) report the following experiment: each participant receives a questionnaire asking him to make two choices, the first from $\{a, b\}$ and *the second from* $\{c,d\}$ *:*

a. *A sure profit of* \$240*.*

b. A lottery between a profit of \$1000 *with probability* 25% *and* 0 *with probability 75%.*

c. A sure loss of \$750*.*

d. A lottery between a loss of \$1000 *with probability* 75% *and* 0 *with probability 25%.*

The participant will receive the sum of the outcomes of the two lotteries he chooses. 73% of the participants chose the combination a and d. Is their behavior sensible?

The combination of a+d is the lottery $0.75[-760$] + 0.25[+240]$ whereas the combination b+c is the lottery $0.75[-750$] + 0.25[+250]$ which first order stochastically dominates $a + d$.

Problem C3 (NYU 2007)

Identify a professor's lifetime with the interval [0,1]. There are $K + 1$ academic *ranks,* 0,..,*K. All professors start at rank* 0 *and eventually reach rank K. Define a career as a sequence* $t = (t_1, \ldots, t_K)$ *where* $t_0 = 0 \le t_1 \le t_2 \le \ldots \le t_K \le 1$ *with the interpretation that* t_k *is the time it takes to get the k'th promotion. (Note that a professor can receive multiple promotions at the same time.) Denote by the professor's preferences on the set of all possible careers.*

For any $\epsilon > 0$ *and for any career t* such that $t_K \leq 1 - \epsilon$, define $t + \epsilon$ to be the *career* $(t + \epsilon)_k = t_k + \epsilon$ **for all k** (i.e. all promotions are delayed by ϵ).

Following are two properties of the professor's preferences:

Monotonicity: For any two careers t and *s*, if $t_k \leq s_k$ for all *k* then $t \geq s$ and if $t_k < s_k$ for all *k*, then $t > s$.

Invariance: For every $\epsilon > 0$ *and every two careers t and s* for which $t + \epsilon$ *and s* + ϵ are well defined, $t \geq s$ iff $t + \epsilon \geq s + \epsilon$.

1. *Formulate the set L of careers in which a professor receives all K promotions at the same time. Show that if satisfies continuity and monotonicity, then for every career t there is a career* $s \in L$ *such that* $s \sim t$ *.*

 $L = \{s \mid s = (\alpha, \ldots, \alpha) \text{ for some } \alpha \in [0,1]\}.$

By monotonicity, $(0,\ldots,0) \ge t \ge (1,\ldots,1)$. By continuity, there exists a bundle *s* on the interval connecting $(0, \ldots, 0)$ and $(1, \ldots, 1)$ such that $s \sim t$. Clearly, $s \in L$ since it is on the main diagonal.

2. *Show that any preference which is represented by the function* $U(t) = -\sum \Delta_k t_k$ *(for some* $\Delta_k > 0$ *) satisfies Monotonicity, Invariance and Continuity.*

MON: If $t_k \leq s_k$ for all k, then $-\Delta_k t_k \geq -\Delta_k s_k$ for all k. Consequently,

 $U(t) = \sum -\Delta_k t_k \geq \sum -\Delta_k s_k = U(s)$, and thus $t \geq s$, and analogously in the strict case. INV: *t s* −∑Δ*ktk* ≥ −∑Δ*ksk* −∑Δ*ktk* − ∑Δ*^k* ≥ −∑Δ*ksk* − ∑Δ*^k* $\Leftrightarrow -\sum \Delta_k(t_k+\epsilon) \geq -\sum \Delta_k(s_k+\epsilon) \Leftrightarrow t+\epsilon \geq s+\epsilon.$ CON: *U* is continuous, so \geq is continuous.

3. *One professor evaluates a career by the maximum length of time one has to wait for a promotion and the smaller this number the better. Show that these preferences cannot be represented by the utility function described in (2).*

 $(.2, .8) \sim (.1, .7)$, and thus preferences fail monotonicity.

Problem C4 (NYU 2008)

An economic agent has to choose between projects. The outcome of each project is uncertain. It might yield a failure or one of K "types of success". Thus, each project z can be described by a vector of K non-negative numbers, (z_1, \ldots, z_K) where z_k stands for the probability that the project success will be of type k .

Let Z ⊂ *^K be the set of feasible projects. Assume Z is compact, convex and satsifies "free disposal".*

The decision maker is an Expected Utility maximizer.

Denote by uk the vNM utility from the k–th type of success, and attach 0 *to failure. Thus the decision maker chooses a project (vector) z* ∈ *Z in order to maximize* $\sum z_k u_k$ *.*

1. *First, formalize the decision maker's problem. Then, formalize (and prove) the claim: If the decision maker suddenly values type k success higher than before, he would choose a project assigning a higher probability to k.*

The DM solves:

$$
max_{z \in Z} z \cdot u = max_{z \in Z} \sum_{k=1}^{K} z_k u_k
$$

Claim: Let $u'_i = u_i$ for every $i \neq k$, $u'_k > u_k$. Then $z_k^* \leq z_k^*$ (where z^* and $z^{*'}$ are the solutions for *u* and *u*′ respectively.

Proof: Equivalent to the proof of "The Law of Demand (or Supply)". $(z^* - z^{*}) \cdot [u - u'] = [z^* - z^{*'}] \cdot u + [z^{*'} - z^*] \cdot u' \ge 0$, since $z^* \cdot u \ge z^{*'} \cdot u$, and *z*∗′ • $u' \ge z$ ∗ • u' . Since $u - u' = (0, ..., 0, u_k - u'_k, 0, ..., 0) < 0$, then $z_k^* - z_k^{*'} \le 0$ QED.

2. *Apparently, the decision maker realizes that there is an additional uncertainty. The world may go "one way or another". With probability* α the vNM *utility of the k'th type of success will be* u_k *and with probability* $1 - \alpha$ *it will* v_k *. Failure remains* 0 *in both contingencies.*

First, formalize the decision maker's new problem. Then, formalize (and prove) the claim: Even if the decision maker would obtain the same expected utility, would he have known in advance the direction of the world, the existence of uncertainty makes him (at least weakly) less happy.

The DM solves:

$$
max_{z\in Z} \cdot [\alpha u + (1-\alpha)v] = max_{z\in Z} \sum_{k=1}^{K} z_k [\alpha u_k + (1-\alpha)v_k]
$$

Claim: The maximal expected utility in the uncertain world is weakly less than the maximal expected utility when the direction of the world is known.

Proof: Denote the DM's chosen projects under the two seperate directions (with vNM utility *u* or *v*) as *zu* and *zv*.

Then

$$
\max_{z \in Z} \mathbf{C} \cdot [\alpha u + (1 - \alpha)v] \le \alpha [\max_{z' \in Z} z' \cdot u] + (1 - \alpha) [\max_{z' \in Z} z' \cdot v] = \alpha z^u \cdot u + (1 - \alpha) z^v \cdot v
$$

Even if $z^u \cdot u = z^v \cdot v = E$, we get $\max_{z \in Z} \mathbf{C} \cdot [\alpha u + (1 - \alpha)v] \le E$.

Problem C5 (NYU 2009)

For any non negative integer n and a number $p \in [0,1]$ let (n,p) be the lottery *which gets the prize* \$*n with probability p and* \$0 *with probability* 1 − *p. Let us call those lotteries "simple lotteries". Consider preference relations on the space of simple lotteries.*

We say that such a preference relation satisfies Independence if $p \geq q$ *iff* $ap \oplus (1 - a)r > aa \oplus (1 - a)r$ for any $a > 0$, and any simple lotteries p, a, r for which *the compound lotteries are also simple lotteries.*

Consider a preference relation satisfying the Independence axiom, strictly monotonic in money and continuous in p. Show that:

1. (n, p) *is monotonic in p* for $n > 0$, *i.e. for all* $p > p'$ $(n, p) > (n, p')$ Observation 1: By monotonicity, $(n,1)$ $>$ $(m,1)$ for all $m < n$. Observation 2: For all $n, (n,0) \sim (0,1)$ since both lotteries give 0 w.p. 1. Proof of Claim:

By observation 1 and 2, $(n,1)$ \times $(n,0)$. By independence $p'/p(n,1) \oplus (1-p'/p)(n,1) \succ p'/p(n,0) \oplus (1-p'/p)(n,1) \Rightarrow (n,1) \succ (n,p'/p).$ Using independence again, $p(n, 1) ⊕ (1-p)(n,0) > p(n,p'/p) ⊕ (1-p)(n,0) ⇒$ (n,p) $>$ (n,p')

2. For all *n* there is a unique $v(n)$ such that $(1,1) \sim (n,1/v(n))$

By observations above for $n > 1$, $(n,1) > (1,1) > (n,0)$. Since (n,p) is continuous, and monotonic in *p*, there exists a unique p_n such that $(1,1) \sim (n,p_n)$. Denote $v(n)$ such that $v(n) = 1/p_n$, and $v(0) = 0$, and naturally $v(1) = 1$

3. It can be represented with the expected utility formula: that is there is an increasing function v such that pvn is a utility function which represents the preference relation.

Claim : For $n > m$, $v(n) > v(m)$

By monotonicity in money, $(n,1)$ $>$ $(m,1)$. By independence

 $(n, 1/v(n)) \sim (1, 1) \succ (m, 1/v(n)) \Rightarrow (m, 1/v(m)) \succ (m, 1/v(n))$. By monotonicity in *p*, $1/\nu(m) > 1/\nu(n)$

Now lets check that $u(n, p) = v(n)p$ represents preferences over these lotteries. Note that, $(n,1/v(n)) \sim (m,1/v(m))$. By independence $(n,v(m)/v(n)q) \sim (m,q)$. Then (n, p) relates to (m, q) like *p* relates to $v(m)/v(n)q$. Thus $(n, p) > (m, q)$ iff $v(n)p > v(m)q$.

D. Social Choice:

Problem D1 (Princeton 2000)

Consider the following social choice problem: a group has n members who must choose from a set containing 3 *elements A*,*B*,*L, where A and B are prizes and L is the lottery which yields each of the prizes A and B with equal probability. Each member has a strict preference over the three alternatives that satisfies vNM assumptions.*

1. *Show that there is a non-dictatorial social welfare function which satisfies the independence of irrelevant alternatives axiom (even the strict version I*[∗]*) and the Pareto axiom (Par).*

Since the preference relations is strict and sastifies vNM assumptions, each of them must be of one of two types

 $A \succ_A L \succ_A B$ or $B \succ_B L \succ_B A$. Let \sim be the preference $A \sim L \sim B$ (which also satsifies the vNM assumptions).

The majority rule assigns the relation

 \succ_A *if* $\{i \mid \succ_i = \succ_A \} > \{i \mid \succ_i = \succ_B \}$ \sim *if* $\{i | \succ_i = \succ_A \} = \{i | \succ_i = \succ_B \}$. \succ_B *if* $\{i \mid \succ_i = \succ_B \} > \{i \mid \succ_i = \succ_A \}$

Clearly, this satisfies the Pareto axiom - if every individual prefers x to v , then so does society, according to the majority rule.

It also satisfies IIA: consider two profiles $\{\succsim_i\}$ and $\{\succsim'_i\}$, and pick two pairs of alternatives (a,b) and (x,y) s.t. $a \succsim_i b$ iff $x \succsim'_i y$. Note that every preference relation in the domain of the SWF corresponds to a distinct preference over any two alternatives. Thus if $a \succ_i b$ iff $x \succ'_i y$ then the relation \succ_{ab} where $a \succ_{ab} b$ is most popular iff the relation and \succ_{xy} where $x \succ_{xy} y$ is most popular.

2. *Reconcile this fact with Arrow's Impossibility Theorem.*

Arrow's impossibility theorem says that if there are at least three alternatives, then the only SWF with unrestricted domain satisfying the Pareto and IIA axioms is the dictatorship. But here the domain is restricted to only two possible preference relations.

Problem D2 (NYU 2009)

We will say that a choice function C is consistent with the majority vetoes a dictator procedure if there are three preference relations \succ_1 , \succ_2 *and* \succ_3 *such that* $c(A)$ *is the* \succ_1 *maximum unless both* \succ_2 *and* \succ_3 *agree on another alternative being the maximum in A.*

1. Show that such a choice function might not be rationalizable.

We will show that the choice function violates property α . Consider the following preferences on *a*,*b*, *c*:

$$
a \succ_1 b \succ_1 c
$$

\n
$$
b \succ_2 a \succ_2 c
$$

\n
$$
c \succ_3 b \succ_3 a
$$

According to these preferences

$$
C(\{a,b,c\}) = a
$$

$$
C(\{a,b\}) = b
$$

2. Show that such a choice function satisfies the following property: If $c(A) = a$, $c(A - \{b\}) = c$ for *b* and *c* different than *a* then $c(B) = c$ for all *B* which contains *c and is a subset of* $A - \{b\}$.

Claim: *a* is \succ_1 maximal in *A*. Assume not for contradiction. Then, *a* must be \succ_2 and \succ_3 maximal in *A*. But since $A - \{b\}$ is a subset of *A*, *a* must be \succ_2 and \succ_3 maximal there, too. But according to the majority veto dictator rule it must also be chosen in $A - \{b\}$ which contradicts $c(A - \{b\}) = c$.

By claim, and $A - \{b\} \subset A$, we know that a is \succ_1 maximal in $A - \{b\}$. Since $c(A - \{b\}) = c$, we know that *c* must be \succ_2 and \succ_3 maximal in $A - \{b\}$. (Otherwise *a* would be chosen in $A - \{b\}$.) But *c* must be maximal \succ_2 and \succ_3 in any *B* which contains *c* and is a subset of $A - \{b\}$. Then *c* must be chosen in any of these subsets.

(c) Show that not all choice functions could be explained by the majority vetoes a dictator procedure.

Any choice function satisfying $C({a,b,c,d}) = a$, $C({a,c,d}) = c$, and $C({a,c}) = a$ violates the property in part (2) and thus cannot be explained by the majority vetoes a dictator procedure.

Problem D3 (Tel Aviv 2009. Inspired by Miller (2007)).

Lately we have been using the term a "reasonable reaction" quite frequently. In this problem we assume that this term is defined according to the opinions of the individuals in the society with regard to the question:"What is a reasonable reaction?".

Assume that in a certain situation, the possible set of reactions is X and the set of individuals in the society is N.

A "reasonability perception" is a non-empty set of possible reactions that are perceived as reasonable.

The social reasonability perception is determined by a function f which attaches a reasonability perception (a non-empty subset of X) to any profile of the individuals' reasonability perception (a vector of non-empty subsets of X).

1. Formalize the following proposition:

Assume that the number of reactions in X is larger than the number of individuals in the society and that f satisfies the following four properties:

A. If in a certain profile all the individuals do not perceive a certain reaction as reasonable, then neither does the society.

B. All the individuals have the same status.

C All the reactions have the same status.

D. Consider two profiles that are different only in one individual's reasonability perception. Any reaction that f determines to be reasonable in the first profile, and regarding which the individual did not change his opinion from reasonable to unreasonable in the second profile, remains reasonable.

Then f determines that a reaction is socially reasonable if and only if at least one of the individuals perceives it as reasonable.

Denote by *Si* the reasonability perception of individual *i*.

Proposition:

Assume $|X| > |N|$. Let *f* be a function that satisfies:

A. $\forall i \in N$. $x \notin S_i \Rightarrow x \notin f(\{S_i\}_{i \in N})$.

B. Let σ be a permutation of *N*. If $\{S_i\}_{i\in N}$ and $\{S_i'\}_{i\in N}$ are two reasonability perception profiles such that for every $i\colon S_i^{'}=S_{\sigma(i)}$ then $f(\{S_i\}_{i\in N})=f(\{S_i^{'}\}_{i\in N})$.

C. Let σ' be a permutation of *X*. If $\{S_i\}_{i\in N}$ and $\{S_i'\}_{i\in N}$ are two reasonability perception profiles such that for every *x* and for every *i* it holds that $x \in$ $S_i \Leftrightarrow \sigma'(x) \in S_i^{'}$, then $x \in f(\{S_i\}_{i \in N}) \Leftrightarrow \sigma'(x) \in f(\{S_i^{'}\}_{i \in N})$.

D. Let $\{S_i\}_{i\in\mathbb{N}}$ and $\{S_i'\}_{i\in\mathbb{N}}$ be two reasonability perception profiles such $S_i^{'}=S_i$ that for any $i \neq j$. Let $x \in f(\langle S_i \rangle_{i \in N})$. If $x \neq S_j$ or $x \in S_j^{'}$, then $x \in f(\langle S_i^{'} \rangle_{i \in N})$.

Then, $x \in f(\langle S_i \rangle_{i \in N}) \Leftrightarrow \exists i \text{ such that } x \in S_i$.

2. Show that all four properties are necessary for the proposition.

1. The fixed function $f(\cdot) = X$ satisfies B, C and D but not A.

2. The function $f(\{S_i\}_{i\in\mathbb{N}}) = S_i$ for some fixed *i* (a dictatorship) satisfies A, C and D but not B.

3. A function which determines that a reaction is reasonable if and only if at least one of the individuals perceives it as such, except for one specific reaction for which it is necessary that two individuals perceive it as reasonable, satisfies A, B

and D but not C.

4. A function which determines as reasonable the reaction(s) that the largest number of individuals perceive as reasonable (the most popular reaction(s)), satisfies A, B and C but not D.

3. Prove the proposition.

 \Rightarrow

Let *f* be a function satisfying A, B, C and D. Let $\{S_i\}_{i\in\mathbb{N}}$ be a reasonability perception profile. Let *y* be a reaction that is perceived as reasonable by at least one individual, denoted by *j*.

Define the profile $\{T_i\}_{i\in\mathbb{N}}$ by arbitrarily assigning one alternative to each individual, with no repetitions (that is $T_i = \{x_i\}$ and $x_i \neq x_k$ for any $i \neq k$), such that $x_i = y$. This is possible since there are more alternatives than individuals.

Claim: In the profile $\{T_i\}_{i\in\mathbb{N}}$, all reactions are determined to be socially reasonable.

Proof: $f(\{T_i\}_{i\in N})$ is non-empty and therefore for some $x, x \in f(\{T_i\}_{i\in N})$. By property A, there is an individual *k* such that $x = x_k \in T_k$. Let $i \neq k$ and let σ be a permutation of *N* that switches between *i* and *k*. Now $\{x_k\} = T_i$ and $\{x_i\} = T_k$ and by property B, $x_k \in f(\{T_{\sigma(i)}\}_{i \in N})$. Let $\sigma^{'}$ be a permutation of X that switches between x_i and x_k . By property C, x_i is now socially reasonable, but in fact we are back to the original profile $\{T_i\}_{i\in\mathbb{N}}$. Therefore, for every *i*, it holds that $x_i \in f(\{T_i\}_{i\in\mathbb{N}})$.

The above claim implies that $y \in f(\langle T_i \rangle_{i \in N})$.

Define ${R_i}_{i \in N}$ such that for every *i*, $R_i = S_i \cup T_i$. One can transform ${T_i}_{i \in N}$ into *Riⁱ*∈*^N* by adding one reaction to one individual at a time. By property D, in each of these stages, the reaction *y* remains socially reasonable and thus $y \in f(\lbrace R_i \rbrace_{i \in N})$.

If for some individual i the reaction that we chose arbitrarily in T_i is not in the original reasonability perception set *Si*, then subtract it from *Ri*. After a finite number of subtractions, we will obtain the original profile $\{S_i\}_{i\in\mathbb{N}}$. Since at no step did we change the status of *y* in any reasonability perception set, $y \in f(\{S_i\}_{i\in N})$ by D.

 \leftarrow

Trivial.

Problem D5 (Tel Aviv 2010)

Let \geq be a preference relation on $Rⁿ$ satisfying the following two properties: *Weak Pareto (WP):* If $x_i \geq y_i$ for all i, then $x = (x_1, \ldots, x_n) \geq y = (y_1, \ldots, y_n)$ and if $x_i > y_i$ for all *i*, then $(x_1, \ldots, x_n) > (y_1, \ldots, y_n)$.

Independence (IIA): Let $a, b, c, d \in \mathbb{R}^n$ be vectors such that in any coordinate $a_i > b_i$, $a_i = b_i$ or $a_i < b_i$ *if and only if* $c_i > d_i$, $c_i = d_i$ or $c_i < d$, accordingly. Then, $a \geq b$ *iff* $c \geq d$ *.*

1. Find a preference relation different from those represented by $u_i(x_1,...,x_n) = x_i$ which satisfies the two properties.

Lexicographic preferences such as: $x > y$ iff $x_{i^*} > y_{i^*}$ where $i^* = \min\{i|x_i \neq y_i\}$.

2. *Show, for* $n = 2$, that there is an *i* such that $a_i > b_i$ implies $a > b$.

Assume that $(4,2) > (2,4)$. By Pareto $(4,2) > (2,0)$.

Also $(4,2)$ $>$ $(2,2)$ since by by Pareto $(2,4) \geq (2,2)$.

Now, consider two vectors $a = (a_1, a_2)$ and $b = (b_1, b_2)$ such that $a_1 > b_1$. By IIA the preference between *a* and *b* when $a_2 < b_2$, $a_2 = b_2$ or $a_2 > b_2$ is the same as between (4,2) and (2,4), (2,2) or (2,0) respectively, namely $a > b$.

3. Provide a "social choice" interpretation for the result in (2). Explain how it differs from Arrow's Impossibility Theorem.

We can interpret a point in $Rⁿ$ as an allocation of a desirable good between *n* individuals. The preferences of all individuals are fixed (each wants as much as possible). The independence property expresses a requirement that the social preference between any two alteratives *a* and *b* is a function of only the *n* comparisons between *ai* and *bi*.

4. Expand (2) for any n.

Let $A \subseteq \{1, ..., n\}$. We say that *A* is *decisive* if whenever for all $i \in A$, $x_i > y_i$ then $x \succ y$.

Let $A \subseteq \{1, \ldots, n\}$. We say that *A* is *almost decisive* if whenever for all $i \in A$, $x_i > y_i$ and for all $i \notin A$, $y_i > x_i$ then $x > y_i$

First, if *A* is almost decisive then it is decisive: By the independence it is enough to look at two vectors *a* and *b* such that $a_i = 3$ and $b_i = 1$ if $i \in A$, and all other a_i and *bj* are either 1 or 3.

Let $c_i = 3$ if $i \in A$ and $c_i = 1$ otherwise and let $d_i = 1$ if $i \in A$ and $d_i = 3$ otherwise.

By the almost decisiveness of *A*, $c > d$. By Pareto $a \geq c$ and $d \geq b$, thus $a > b$. Now let *A* be a decisive set. and let *A*¹ and *A*² be a partition of *A*. We will see

that either A_1 or A_2 is almost decisive.

Assume not. Consider the three vectors:

By *A*'s decisiveness, $c > a$. If A_1 is not almost decisive then $a \ge b$ and if A_2 is not almost decisive then $b \geq c$. A contradiction.

Thus, there is i such that $\{i\}$ is decisive.